

PI 1203199

PCD 0 6 AUG 2004

PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

August 03, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/551,551

FILING DATE: March 09, 2004

RELATED PCT APPLICATION NUMBER: PCT/US04/08700

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



L. Wallace
T. WALLACE
Certifying Officer

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

PATENT APPLICATION SERIAL NO. _____

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE
FEE RECORD SHEET

03/12/2004 HBELETE1 00000072 60551551

01 FC:2005

80.00 DP

PTO-1556
(5/87)



Attorney's Docket No: 046562/273962
Express Mail Label No. EV 390050859 US

PATENT

17513 U.S. PTO
20/551551

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

Mail Stop Provisional Patent Application
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 C.F.R. 1.53(c).

INVENTOR(s)

Name: Matthew Oliver Fraser
Residence: 408 Gablefield Lane
Apex, NC 27502

Name: Steven B. Landau
Residence: 44 Tanglewood Road
Wellesley, MA 02481

TITLE OF THE INVENTION (500 characters maximum)

METHODS FOR TREATING FUNCTIONAL BOWEL DISORDERS USING
 $\alpha 2\delta$ SUBUNIT CALCIUM CHANNEL MODULATORS WITH SMOOTH
MUSCLE MODULATORS

CORRESPONDENCE ADDRESS

Edward R. Ergenzinger, Jr.
Registration No. 47,549
Customer Number 00826

Tel Raleigh Office (919) 862-2200
Fax Raleigh Office (919) 862-2260

ENCLOSED APPLICATION PARTS (check all that apply)

- | | |
|-------------------------------------|---|
| <input checked="" type="checkbox"/> | Specification (Number of Pages <u>141</u>) |
| <input type="checkbox"/> | Drawing(s) (Number of Sheets <u> </u>) |
| <input type="checkbox"/> | Application Data Sheet. See 37 CFR 1.76 |
| <input type="checkbox"/> | CD(s), Number |
| <input type="checkbox"/> | Other (specify) |

METHOD OF PAYMENT OF FILING FEES

- ☒ Applicant claims small entity status
☒ Check or money order is enclosed to cover the filing fee.
☐ The Commissioner is hereby authorized to charge filing fees to Deposit Account No. 16-0605.
☒ Please charge Deposit Account No. 16-0605 for any fee deficiency.

PROVISIONAL FILING FEE AMOUNT(s)

Large Entity \$160.00; Small Entity \$ 80.00

Filing Fee Amount: \$80.00

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

- ☒ No.
☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,

Edward R. Ergenzinger, Jr.

Edward R. Ergenzinger, Jr.

Registration No. 47,549

<p>CUSTOMER NO. 00826 ALSTON & BIRD LLP Bank of America Plaza 101 South Tryon Street, Suite 4000 Charlotte, NC 28280-4000 Tel Raleigh Office (919) 862-2200 Fax Raleigh Office (919) 862-2260</p>	<p>"Express Mail" mailing label number EV 390050859 US Date of Deposit: March 9, 2004</p> <p>I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to: Mail Stop Provisional Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450</p> <p><i>Lynda Jo Pixley</i> Lynda Jo Pixley</p>
---	---

**METHODS FOR TREATING FUNCTIONAL BOWEL DISORDERS USING $\alpha_2\delta$
SUBUNIT CALCIUM CHANNEL MODULATORS WITH SMOOTH MUSCLE
MODULATORS**

FIELD OF THE INVENTION

The invention relates to methods for treating functional bowel disorders, particularly Irritable Bowel Syndrome, using smooth muscle modulators and $\alpha_2\delta$ subunit calcium channel modulators.

BACKGROUND OF THE INVENTION

Functional Bowel Disorders (FBDs) refer to disorders or diseases where the primary abnormality is an altered physiological function of the bowels, rather than an identifiable structural or biochemical cause. While the term bowel is commonly defined as the small and large intestines, the phrase "functional bowel disorder" is generally held to encompass disorders or diseases related to the mid or lower gastrointestinal tract, which includes the stomach as well as the intestines. Specific FBD's are Irritable Bowel Syndrome (IBS), functional abdominal bloating, functional constipation, functional diarrhea, and unspecified functional bowel disorder (see, Drossman D. *et al.* [Eds.], *Functional GI Disorders: Diagnosis, Pathophysiology, Treatment, A Multinational Consensus*, 1994). Additional gastrointestinal disorders are also viewed as being encompassed by the FBD designation, including functional dyspepsia (see, Camilleri, M., *Functional Bowel Disease: Roles of Sensation and Motility* (64th Annual Meeting of the Swiss Society of Gastroenterology and Hepatology), *Schweiz Med. Wochenschr*, 2000, 130:1772-81), and functional abdominal pain syndrome (see CME series No. 13, available online at the website: medicine.org.hk/hkducme/bulletin/200307/27.pdf).

FBDs are diagnosed by characteristic symptoms being present for at least 12 weeks during the preceding 12 months in the absence of a structural or biochemical explanation (see, Thompson *et al.*, *Gut*, 45 (Suppl. II):II43-II47 (1999)). Individual FBDs (IBS, functional abdominal bloating, functional constipation, and functional

diarrhea) are distinguished by symptom-based diagnostic criteria. Unspecified FBD presents as a bowel disorder lacking the specific criteria for the other FBDs.

These specified, and unspecified, FBDs are known to affect a large number of individuals. IBS alone is estimated to affect approximately 10-20% of the general population. IBS is the most common disease diagnosed by gastroenterologists and one of the most common disorders seen by all physicians. Since IBS has no characteristic pathophysiological abnormality, the diagnosis is mainly based on symptom analysis, such as by the Manning criteria (1978) or the Rome I (1989) and Rome II (1999) criteria. Generally, symptoms associated with IBS include abdominal pain, abnormal stool frequency, abnormal stool form, abnormal stool passage, mucorrhea, and bloating or feeling of abdominal distension. A diagnosis of IBS is generally supported through cumulative presence of the symptoms described above.

Treatment options for IBS generally encompass multiple approaches that are customized to the patient depending upon the severity of the symptoms. Patients diagnosed with mild IBS symptoms are often counseled about managing stress and making diet and lifestyle changes. Patients diagnosed with moderate IBS are similarly counseled with the added recommendation of fiber supplements. Depending on the symptoms, moderate IBS patients can also be advised to use antidiarrheals, laxatives, or anticholinergic agents. Typical antidiarrheals include loperamide (Imodium[®]), attapulgite (Kaopectate[®]), and diphenoxylate (Lotomil[®]). Typical laxatives include bisacodyl (Dulcolax[®]), senna (Senokot[®]), polyethylene 3350 (Miralax[™]), and bulk-forming fiber laxatives, such as psyllium (Metamucil[®]), calcium polycarbophil (Equalactin[®]), methylcellulose (Citrucel[®]), and fructan (Fiber Choice[®]). An example of an anticholinergic used in treating IBS is dicyclomine (Bentyl[®]).

Patients diagnosed with severe IBS may receive, in addition to the above, treatment with antidepressants, including tricyclics and selective serotonin reuptake inhibitors (SSRIs). Severe IBS may also be treated with medication having 5-HT₃ activity, such as alosetron (Lotronex[®]), or 5-HT₄ activity, such as tegaserod (Zelnorm[®]).

Tricyclic antidepressants, such as amitriptyline (Amitril[®] or Elavil[®]), imipramine (Tofranil[®]), and doxepin (Adapin[®]), are used in treating IBS symptoms because of the anticholinergic and analgesic properties they exhibit independent of their psychotropic

effects. It has been reported that the anticholinergic and analgesic effects of the tricyclic antidepressants on the gastrointestinal tract can benefit patients with pain predominant IBS and increased bowel frequency (see, Clousse, R.R., *Dig. Dis. Sci.*, 39: 2352-2363 (1994)).

Given the variable dynamics of IBS, problems associated with treatment, especially long-term treatment, abound. Identifying diet or lifestyle patterns that promote IBS can be difficult to accomplish, and such changes can have limited effectiveness. Antidiarrheal medications often cause side effects, such as bloating, cramping, and constipation. Laxative dependency is of itself problematic and can cause fluctuating bowel habits that lead to social difficulties. Further, many patients are also reluctant to use drugs associated with psychiatric indications, such as depression.

Because existing therapies and treatments for functional bowel disorders, especially IBS, are associated with limitations, as described above, new therapies and treatments are therefore desirable.

SUMMARY OF THE INVENTION

Compositions and methods for treating functional bowel disorders, particularly IBS, are provided. Compositions of the invention comprise $\alpha_2\delta$ subunit calcium channel modulators in combination with one or more compounds with smooth muscle modulatory effects. According to the present invention, $\alpha_2\delta$ subunit calcium channel modulators include GABA analogs (e.g., gabapentin and pregabalin), fused bicyclic or tricyclic amino acid analogs of gabapentin, and amino acid compounds. Compounds with smooth muscle modulatory effects include antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors. Compositions of the invention include combinations of the aforementioned compounds as well as pharmaceutically acceptable, pharmacologically active acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

The compositions are administered in therapeutically effective amounts to a patient in need thereof for treating functional bowel disorders. It is recognized that the compositions may be administered by any means of administration as long as an effective amount for the treatment of symptoms associated with functional bowel disorders. The

compositions may be formulated, for example, for sustained, continuous, or as-needed administration.

DETAILED DESCRIPTION OF THE INVENTION

Overview and Definitions

The present invention provides compositions and methods for treating functional bowel disorders. The compositions comprise a therapeutically effective dose of a compound with smooth muscle modulatory effects in combination with an $\alpha_2\delta$ subunit calcium channel modulator, such as gabapentin or pregabalin. Compounds with smooth muscle modulatory effects include, but are not limited to, antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors. The methods are accomplished by administering, for example, a compound with smooth muscle modulatory effects, such as oxybutynin, in combination with an $\alpha_2\delta$ subunit calcium channel modulator and/or another compound that interacts with $\alpha_2\delta$ subunit-containing calcium channels, such as gabapentin or pregabalin. For these methods, various compositions and formulations that contain quantities of a compound with smooth muscle modulatory effects in combination with an $\alpha_2\delta$ subunit calcium channel modulator and/or other compounds that interact with $\alpha_2\delta$ subunit-containing calcium channels are encompassed.

It is to be understood that the inventions are not to be limited to the specific embodiments disclosed and that modifications and other embodiments are intended to be included within the scope of the appended claims. Although specific terms are employed herein, they are used in a generic and descriptive sense only and not for purposes of limitation.

It must be noted that as used in this specification and the appended embodiments, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "an active agent" or "a pharmacologically active agent" includes a single active agent as well as two or more different active agents in combination, reference to "a carrier" includes mixtures of two or more carriers as well as a single carrier, and the like.

By "non-painful" is intended sensations or symptoms including mild or general discomfort that a patient subjectively describes as not producing or resulting in pain.

By "painful" is intended sensations or symptoms that a patient subjectively describes as producing or resulting in pain.

The terms "active agent" and "pharmacologically active agent" are used interchangeably herein to refer to a chemical compound that induces a desired effect, i.e., in this case, treatment of functional bowel disorders. The primary active agents herein are $\alpha_2\delta$ subunit calcium channel modulators and/or smooth muscle relaxants. The present invention comprises a combination therapy wherein an $\alpha_2\delta$ subunit calcium channel modulator is administered with one or more smooth muscle modulator. Such combination therapy may be carried out by administration of the different active agents in a single composition, by concurrent administration of the different active agents in different compositions, or by sequential administration of the different active agents. Included are derivatives and analogs of those compounds or classes of compounds specifically mentioned that also induce the desired effect.

The term " $\alpha_2\delta$ subunit calcium channel modulator" as used herein refers to an agent that is capable of interacting with the $\alpha_2\delta$ subunit of a calcium channel, including a binding event, including subtypes of the $\alpha_2\delta$ calcium channel subunit as disclosed in Klugbauer et al. (1999) *J. Neurosci.* 19: 684-691, to produce a physiological effect, such as opening, closing, blocking, up-regulating functional expression, down-regulating functional expression, or desensitization, of the channel. Unless otherwise indicated, the term " $\alpha_2\delta$ subunit calcium channel modulator" is intended to include GABA analogs including gabapentin and pregabalin,, fused bicyclic or tricyclic amino acid analogs of gabapentin, amino acid compounds, and other compounds that interact with the $\alpha_2\delta$ calcium channel subunit as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

The term "peptidomimetic" is used in its conventional sense to refer to a molecule that mimics the biological activity of a peptide but is no longer peptidic in chemical nature, including molecules that lack amide bonds between amino acids, as well

as pseudo-peptides, semi-peptides and peptoids. Peptidomimetics according to this invention provide a spatial arrangement of reactive chemical moieties that closely resembles the three-dimensional arrangement of active groups in the peptide on which the peptidomimetic is based. As a result of this similar active-site geometry, the peptidomimetic has effects on biological systems that are similar to the biological activity of the peptide.

The term "smooth muscle modulator" as used herein refers to any compound that inhibits or blocks the contraction of smooth muscles, including but not limited to antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors. Smooth muscle modulators can be "direct" (also known as "musculotropic") or "indirect" (also known as "neurotropic"). "Direct smooth muscle modulators" are smooth muscle modulators that act by inhibiting or blocking contractile mechanisms within smooth muscle, including but not limited to modification of the interaction between actin and myosin. "Indirect smooth muscle modulators" are smooth muscle modulators that act by inhibiting or blocking neurotransmission that results in the contraction of smooth muscle, including but not limited to blockade of presynaptic facilitation of acetylcholine release at the axon terminal of motor neurons terminating in smooth muscle.

The term "anticholinergic agent" as used herein refers to any acetylcholine receptor antagonist, including antagonists of nicotinic and/or muscarinic acetylcholine receptors. The term "antinicotinic agent" as used herein is intended any nicotinic acetylcholine receptor antagonist. The term "antimuscarinic agent" as used herein is intended any muscarinic acetylcholine receptor antagonist. Unless otherwise indicated, the terms "anticholinergic agent," "antinicotinic agent," and "antimuscarinic agent" are intended to include anticholinergic, antinicotinic, and antimuscarinic agents as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any acids, salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

The term " β_3 adrenergic agonist" is used in its conventional sense to refer to a compound that binds to and agonizes β_3 adrenergic receptors. Unless otherwise

indicated, the term "β3 adrenergic agonist" is intended to include β3 adrenergic agonist agents as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any acids, salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

The term "spasmolytic" (also known as "antispasmodic") is used in its conventional sense to refer to a compound that relieves or prevents muscle spasms, especially of smooth muscle. Unless otherwise indicated, the term "spasmolytic" is intended to include spasmolytic agents as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

The term "neurokinin receptor antagonist" is used in its conventional sense to refer to a compound that binds to and antagonizes neurokinin receptors. Unless otherwise indicated, the term "neurokinin receptor antagonist" is intended to include neurokinin receptor antagonist agents as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any acids, salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

The term "bradykinin receptor antagonist" is used in its conventional sense to refer to a compound that binds to and antagonizes bradykinin receptors. Unless otherwise indicated, the term "bradykinin receptor antagonist" is intended to include bradykinin receptor antagonist agents as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any acids, salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

The term "nitric oxide donor" is used in its conventional sense to refer to a compound that releases free nitric oxide when administered to a patient. Unless

otherwise indicated, the term "nitric oxide donor" is intended to include nitric oxide donor agents as disclosed further herein, as well as acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof. Further, it is understood that any acids salts, esters, amides, prodrugs, active metabolites or other derivatives are pharmaceutically acceptable as well as pharmacologically active.

"Constipation" is used in its conventional sense to mean infrequent or difficult evacuation of feces.

"Diarrhea" is used in its conventional sense to mean a frequent and generally profuse discharge of loose or fluid evacuations from the intestines without straining.

The terms "treating" and "treatment" as used herein refer to relieving the symptoms or other clinically observed sequelae for clinically diagnosed disorders as described herein, including functional bowel disorders, such as IBS.

By an "effective" amount or a "therapeutically effective amount" of a drug or pharmacologically active agent is meant a nontoxic but sufficient amount of the drug or agent to provide the desired effect, i.e., relieving the symptoms associated with functional bowel disorders, as explained above. It is recognized that the effective amount of a drug or pharmacologically active agent will vary depending on the route of administration, the selected compound, and the species to which the drug or pharmacologically active agent is administered, as well as the age, weight, and sex of the individual to which the drug or pharmacologically active agent is administered. It is also recognized that one of skill in the art will determine appropriate effective amounts by taking into account such factors as metabolism, bioavailability, and other factors that affect plasma levels of a drug or pharmacologically active agent following administration within the unit dose ranges disclosed further herein for different routes of administration.

By "pharmaceutically acceptable," such as in the recitation of a "pharmaceutically acceptable carrier," or a "pharmaceutically acceptable acid addition salt," is meant a material that is not biologically or otherwise undesirable, i.e., the material may be incorporated into a pharmaceutical composition administered to a patient without causing any undesirable biological effects or interacting in a deleterious manner with any of the other components of the composition in which it is contained. "Pharmacologically active" (or simply "active") as in a "pharmacologically active" derivative or metabolite,

refers to a derivative or metabolite having the same type of pharmacological activity as the parent compound. When the term "pharmaceutically acceptable" is used to refer to a derivative (e.g., a salt or an analog) of an active agent, it is to be understood that the compound is pharmacologically active as well, i.e., therapeutically effective for treating functional bowel disorders as defined herein.

By "continuous" dosing is meant the chronic administration of a selected active agent.

By "as-needed" dosing, also known as "*pro re nata*" "prn" dosing, and "on demand" dosing or administration is meant the administration of a single dose of the active agent at some time prior to commencement of an activity wherein suppression of the painful and non-painful symptoms of a functional bowel disorder, particularly IBS in normal and spinal cord injured patients, would be desirable. Administration can be immediately prior to such an activity, including about 0 minutes, about 10 minutes, about 20 minutes, about 30 minutes, about 1 hour, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, or about 10 hours prior to such an activity, depending on the formulation.

By "short-term" is intended any period of time up to and including about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, about 1 hour, about 40 minutes, about 20 minutes, or about 10 minutes after drug administration.

By "rapid-offset" is intended any period of time up to and including about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, about 1 hour, about 40 minutes, about 20 minutes, or about 10 minutes after drug administration.

The term "controlled release" is intended to refer to any drug-containing formulation in which release of the drug is not immediate, i.e., with a "controlled release" formulation, oral administration does not result in immediate release of the drug into an absorption pool. The term is used interchangeably with "non-immediate release" as defined in Remington: The Science and Practice of Pharmacy, Twentieth Ed. (Philadelphia, Pa.: Lippincott Williams & Wilkins, 2000).

The "absorption pool" represents a solution of the drug administered at a particular absorption site, and k_r , k_a , and k_e are first-order rate constants for: 1) release of the drug from the formulation; 2) absorption; and 3) elimination, respectively. For immediate release dosage forms, the rate constant for drug release k_r is far greater than the absorption rate constant k_a . For controlled release formulations, the opposite is true, i.e., $k_r \ll k_a$, such that the rate of release of drug from the dosage form is the rate-limiting step in the delivery of the drug to the target area. The term "controlled release" as used herein includes any nonimmediate release formulation, including but not limited to sustained release, delayed release and pulsatile release formulations.

The term "sustained release" is used in its conventional sense to refer to a drug formulation that provides for gradual release of a drug over an extended period of time, and that preferably, although not necessarily, results in substantially constant blood levels of a drug over an extended time period such as up to about 72 hours, about 66 hours, about 60 hours, about 54 hours, about 48 hours, about 42 hours, about 36 hours, about 30 hours, about 24 hours, about 18 hours, about 12 hours, about 10 hours, about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, or about 1 hour after drug administration.

The term "delayed release" is used in its conventional sense to refer to a drug formulation that provides for an initial release of the drug after some delay following drug administration and that preferably, although not necessarily, includes a delay of up to about 10 minutes, about 20 minutes, about 30 minutes, about 1 hour, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, or about 12 hours.

The term "pulsatile release" is used in its conventional sense to refer to a drug formulation that provides release of the drug in such a way as to produce pulsed plasma profiles of the drug after drug administration. The term "immediate release" is used in its conventional sense to refer to a drug formulation that provides for release of the drug immediately after drug administration.

The term "immediate release" is used in its conventional sense to refer to a drug formulation that provides for release of the drug immediately after drug administration.

By the term "transdermal" drug delivery is meant delivery by passage of a drug through the skin or mucosal tissue and into the bloodstream.

The term "topical administration" is used in its conventional sense to mean delivery of a topical drug or pharmacologically active agent to the skin or mucosa.

The term "oral administration" is used in its conventional sense to mean delivery of a drug through the mouth and ingestion through the stomach and digestive tract.

The term "inhalation administration" is used in its conventional sense to mean delivery of an aerosolized form of the drug by passage through the nose or mouth during inhalation and passage of the drug through the walls of the lungs.

The term "intravesical administration" is used in its conventional sense to mean delivery of a drug directly into the bladder.

By the term "parenteral" drug delivery is meant delivery by passage of a drug into the blood stream without first having to pass through the alimentary canal, or digestive tract. Parenteral drug delivery may be "subcutaneous," referring to delivery of a drug by administration under the skin. Another form of parenteral drug delivery is "intramuscular," referring to delivery of a drug by administration into muscle tissue. Another form of parenteral drug delivery is "intradermal," referring to delivery of a drug by administration into the skin. An additional form of parenteral drug delivery is "intravenous," referring to delivery of a drug by administration into a vein. An additional form of parenteral drug delivery is "intra-arterial," referring to delivery of a drug by administration into an artery. Another form of parenteral drug delivery is "transdermal," referring to delivery of a drug by passage of the drug through the skin and into the bloodstream. Another form of parenteral drug delivery is "intrathecal," referring to delivery of a drug directly into the into the intrathecal space (where fluid flows around the spinal cord).

Still another form of parenteral drug delivery is "transmucosal," referring to administration of a drug to the mucosal surface of an individual so that the drug passes through the mucosal tissue and into the individual's blood stream. Transmucosal drug delivery may be "buccal" or "transbuccal," referring to delivery of a drug by passage through an individual's buccal mucosa and into the bloodstream. Another form of transmucosal drug delivery herein is "lingual" drug delivery, which refers to delivery of a

drug by passage of a drug through an individual's lingual mucosa and into the bloodstream. Another form of transmucosal drug delivery herein is "sublingual" drug delivery, which refers to delivery of a drug by passage of a drug through an individual's sublingual mucosa and into the bloodstream. Another form of transmucosal drug delivery is "nasal" or "intranasal" drug delivery, referring to delivery of a drug through an individual's nasal mucosa and into the bloodstream. An additional form of transmucosal drug delivery herein is "rectal" or "transrectal" drug delivery, referring to delivery of a drug by passage of a drug through an individual's rectal mucosa and into the bloodstream. Another form of transmucosal drug delivery is "urethral" or "transurethral" delivery, referring to delivery of the drug into the urethra such that the drug contacts and passes through the wall of the urethra. An additional form of transmucosal drug delivery is "vaginal" or "transvaginal" delivery, referring to delivery of a drug by passage of a drug through an individual's vaginal mucosa and into the bloodstream. An additional form of transmucosal drug delivery is "perivaginal" delivery, referring to delivery of a drug through the vaginolabial tissue into the bloodstream.

In order to carry out the method of the invention, a selected active agent is administered to a patient suffering from a functional bowel disorder. A therapeutically effective amount of the active agent may be administered orally, intravenously, subcutaneously, transmucosally (including buccally, sublingually, transurethrally, and rectally), topically, transdermally, by inhalation, intravesically, intrathecally or using any other route of administration.

Functional Bowel Disorders

Functional Bowel Disorders (FBDs) are generally defined as collections of functional symptoms that are attributable to the mid or lower gastrointestinal tract. A FBD is diagnosed by characteristic symptoms for at least 12 weeks during the preceding 12 months, and such symptoms include abdominal pain, bloating, distention, and various symptoms of disordered defecation. The FBDs are commonly divided into symptom specific disorders including functional diarrhea, functional constipation, functional abdominal bloating, Irritable Bowel Syndrome (IBS), which often encompasses symptoms present in each of the other individual disorders, and unspecified functional

bowel disorder. Additional disorders often classified as FBDs include functional dyspepsia and functional abdominal pain syndrome.

Irritable Bowel Syndrome

IBS is a biopsychosocial disorder with variable symptoms wherein a disturbance in the interaction between intestinal motility and sensation, the brain, and the autonomic nervous system produces the syndrome. IBS is characterized by a group of symptoms where abdominal pain or discomfort is associated with a change in bowel pattern, such as loose stool, more frequent bowel movements, diarrhea, and/or constipation, in the absence of demonstrable organic pathology.

As IBS presents no specific motility or structural correlates, it remains a clinically defined illness defined by either the Manning or Rome II Criteria. The Manning criteria were originally established in 1978 to distinguish IBS from organic bowel disease (Manning A P, Thompson W G, Heaton K W, Morris A F. Towards a positive diagnosis of the irritable bowel. Br. Med. J. 1978; 2:653-4). The criteria are:

1. Onset of pain associated with more frequent bowel movements
2. Onset of pain associated with looser bowel movements
3. Pain relieved by defecation
4. Visible abdominal bloating
5. Subjective sensation of incomplete evacuation more than 25% of the time
6. Mucorrhea more than 25% of the time

The Rome II criteria for IBS include reports for at least 12 weeks in the preceding 12 months, which need not be consecutive, of abdominal pain or discomfort that has two of three features:

1. Relieved with defecation; and/or
2. Onset associated with a change in the frequency of stools; and/or
3. Onset associated with a change in form (appearance) of stool.

Other symptoms, such as abnormal stool frequency, abnormal stool form, abnormal stool passage, passage of mucus, and/or bloating or feeling of abdominal distension, cumulatively support the diagnosis of IBS.

Subjects with IBS exhibit visceral hypersensitivity, the presence of which behavioral studies have shown is the most consistent abnormality in IBS. For example, patients and controls were evaluated for their pain thresholds in response to progressive distension of the sigmoid colon induced by a balloon. At the same volume of distension, the patients reported higher pain scores compared to controls. This finding has been reproduced in many studies and with the introduction of the barostat, a computerized distension device, the distension procedures have been standardized. Two concepts of visceral hypersensitivity, hyperalgesia and allodynia, have been introduced. More specifically, hyperalgesia refers to the situation in which normal visceral sensations are experienced at lower intraluminal volumes. While for a finding of allodynia, pain or discomfort is experienced at volumes usually producing normal internal sensations (see, for example, Mayer E.A. and Gebhart, G.F., *Basic and Clinical Aspects of Chronic Abdominal Pain*, Vol 9, 1 ed. Amsterdam: Elsevier, 1993:3-28).

As such, IBS is a functional bowel disorder in which abdominal pain or discomfort is associated with defecation or a change in bowel habit. Therefore, IBS has elements of an intestinal motility disorder, a visceral sensation disorder, and a central nervous disorder. While the symptoms of IBS have a physiological basis, no physiological mechanism unique to IBS has been identified. In some cases, the same mechanisms that cause occasional abdominal discomfort in healthy individuals operate to produce the symptoms of IBS. The symptoms of IBS are therefore a product of quantitative differences in the motor reactivity of the intestinal tract, and increased sensitivity to stimuli or spontaneous contractions.

In addition to the above diagnosis tools, IBS patients can be categorized according to symptoms and severity. Chronic diarrhea which is associated with abdominal pain and which is not attributable to an organic cause is referred to as "irritable bowel syndrome with a diarrhea predominance" or diarrhea predominant IBS (Hasler et al., 1995, In: *Textbook of Gastroenterology*, 2nd ed., Yamada, Ed., J. B. Lippincott Co., Philadelphia, pp. 1832-1855). Patients exhibit diarrhea predominant IBS if their usual bowel movement frequency is more than three times per day, or if their usual form of stool is loose and not hard, or if they frequently feel the sense of urgency and do not strain at the stools. Patients exhibit constipation predominant IBS if their

usual bowel movement frequency is less than three times per week, or if their usual form of stool is hard and not loose, or if they often strain at the stools and do not frequently feel the sense of urgency. Patient not described as exhibiting diarrhea predominant IBS or constipation predominant IBS can be termed as exhibiting non-specific or alternating constipation/diarrhea IBS.

Functional Abdominal Bloating

Functional abdominal bloating, or abdominal distention, is characterized by a feeling of fullness or bloating throughout the abdomen in the absence of further symptoms associated with another functional gastrointestinal disorder. Functional abdominal bloating is often considered co-morbid with IBS, but studies indicate functional abdominal bloating alone occurs in about 15% of community-based populations, usually with a female predominance.

Functional abdominal bloating is often absent during waking hours and worsens throughout the day. Diagnostic criteria include at least 12 weeks, which need not be consecutive, in the preceding 12 months of: (1) feeling of abdominal fullness, bloating, or visible distension; and (2) insufficient criteria for a diagnosis of functional dyspepsia, IBS, or other functional disorder.

Functional Constipation

Functional constipation is generally characterized by infrequent bowel movements, passage of hard stools, difficulty in passing stools, and seemingly incomplete defecation. While not wishing to be bound by theory, functional constipation may arise from motility problems, including a decrease in the number of high amplitude propagating contractions in the large intestine. Diagnostic criteria include: at least 12 weeks, which need not be consecutive, in the preceding 12 months of two or more of: (1) straining in greater than one-fourth of defecations; (2) lumpy or hard stools in greater than one-fourth of defecations; (3) sensation of incomplete evacuation in greater than one-fourth of defecations; (4) sensation of anorectal obstruction/blockade in greater than one-fourth of defecations; manual maneuvers (e.g. digital evacuation, support of the

pelvic floor) to facilitate greater than one-fourth of defecations; and/or (6) less than three defecations per week.

Functional Diarrhea

Functional diarrhea usually presents as frequent, loose, or watery stools and a subjective sense of urgency, often without the presence of pain. Without wishing to be bound by theory, one cause of functional diarrhea is believed to be an excessive number of high amplitude propagating contractions, which reduce the amount of time food residues remain in the large intestine for water to be reabsorbed. Diagnostic criteria include: at least 12 weeks, which need not be consecutive, in the preceding 12 months of: (1) liquid (mushy) or watery stool; (2) present greater than three-fourths of the time; and (3) no abdominal pain.

Functional Abdominal Pain Syndrome

Functional Abdominal Pain Syndrome (FAPS) is also known as chronic idiopathic abdominal pain or chronic functional abdominal pain. These terms are generally used to describe pain for at least six months that is poorly related to bowel function and is associated with some loss of daily activities. Diagnostic criteria for FAPS include at least six months of: (1) continuous or nearly continuous abdominal pain; and (2) no or only occasional relation of pain with physiological events (e.g. eating, defecation, menses); and (3) some loss of daily functioning; and (4) the pain is not feigned; and (5) insufficient criteria exists for diagnosing other functional gastrointestinal disorders that would explain the abdominal pain.

Functional Dyspepsia

Functional dyspepsia is a functional bowel disorder in which chronic or recurrent symptoms are centered in the upper abdomen without presence of other known disease, such as infection, inflammation, or ulcer. Symptoms include pain and discomfort, which is meant to represent other symptoms, such as early satiety, nausea, vomiting, or bloating. There are two primary motor dysfunctions that can be described in relation to functional dyspepsia. First, more than 30% of adults with functional dyspepsia (or non-ulcer

dyspepsia) have impaired gastric emptying. Second, impaired gastric accommodation is also frequent.

Peripheral vs. Central Effects

The mammalian nervous system comprises a central nervous system (CNS, comprising the brain and spinal cord) and a peripheral nervous system (PNS, comprising sympathetic, parasympathetic, sensory, motor, and enteric neurons outside of the brain and spinal cord). Where an active agent according to the present invention is intended to act centrally (i.e., exert its effects via action on neurons in the CNS), the active agent must either be administered directly into the CNS or be capable of bypassing or crossing the blood-brain barrier. The blood-brain barrier is a capillary wall structure that effectively screens out all but selected categories of substances present in the blood, preventing their passage into the CNS. The unique morphologic characteristics of the brain capillaries that make up the blood-brain barrier are: 1) epithelial-like high resistance tight junctions which literally cement all endothelia of brain capillaries together within the blood-brain barrier regions of the CNS; and 2) scanty pinocytosis or transendothelial channels, which are abundant in endothelia of peripheral organs. Due to the unique characteristics of the blood-brain barrier, hydrophilic drugs and peptides that readily gain access to other tissues in the body are barred from entry into the brain or their rates of entry are very low.

The blood-brain barrier can be bypassed effectively by direct infusion of the active agent into the brain, or by intranasal administration or inhalation of formulations suitable for uptake and retrograde transport of the active agent by olfactory neurons. The most common procedure for administration directly into the CNS is the implantation of a catheter into the ventricular system or intrathecal space. Alternatively, the active agent can be modified to enhance its transport across the blood-brain barrier. This generally requires some solubility of the drug in lipids, or other appropriate modification known to one of skill in the art. For example, the active agent may be truncated, derivatized, latentiated (converted from a hydrophilic drug into a lipid-soluble drug), conjugated to a lipophilic moiety or to a substance that is actively transported across the blood-brain barrier, or modified using standard means known to those skilled in the art.

See, for example, Pardridge, Endocrine Reviews 7: 314-330 (1986) and U.S. Pat. No. 4,801,575.

Where an active agent according to the present invention is intended to act exclusively peripherally (i.e., exert its effects via action either on neurons in the PNS or directly on target tissues), it may be desirable to modify the compounds of the present invention such that they will not pass the blood-brain barrier. The principle of blood-brain barrier permeability can therefore be used to design active agents with selective potency for peripheral targets. Generally, a lipid-insoluble drug will not cross the blood-brain barrier, and will not produce effects on the CNS. A basic drug that acts on the nervous system may be altered to produce a selective peripheral effect by quaternization of the drug, which decreases its lipid solubility and makes it virtually unavailable for transfer to the CNS. For example, the charged antimuscarinic drug methscopolamine bromide has peripheral effects while the uncharged antimuscarinic drug scopolamine acts centrally. One of skill in the art can select and modify active agents of the present invention using well-known standard chemical synthetic techniques to add a lipid impermeable functional group such a quaternary amine, sulfate, carboxylate, phosphate, or sulfonium to prevent transport across the blood-brain barrier. Such modifications are by no means the only way in which active agents of the present invention may be modified to be impermeable to the blood-brain barrier; other well known pharmaceutical techniques exist and would be considered to fall within the scope of the present invention.

Agents

Compounds useful in the present invention include any active agent as defined elsewhere herein. Such active agents include, for example, $\alpha_2\delta$ subunit calcium channel modulators, including GABA analogs (e.g., gabapentin and pregabalin), as described elsewhere herein, as well as smooth muscle modulators, including antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors, as described elsewhere herein.

Voltage gated calcium channels, also known as voltage dependent calcium channels, are multi-subunit membrane-spanning proteins which permit controlled

calcium influx from an extracellular environment into the interior of a cell. Opening and closing (gating) of voltage gated calcium channels is controlled by a voltage sensitive region of the protein containing charged amino acids that move within an electric field. The movement of these charged groups leads to conformational changes in the structure of the channel resulting in conducting (open/activated) or non-conducting (closed/inactivated) states.

Voltage gated calcium channels are present in a variety of tissues and are implicated in several vital processes in animals. Changes in calcium influx into cells mediated through these calcium channels have been implicated in various human diseases such as epilepsy, stroke, brain trauma, Alzheimer's disease, multi-infarct dementia, other classes of dementia, Korsakoff's disease, neuropathy caused by a viral infection of the brain or spinal cord (e.g., human immunodeficiency viruses, etc.), amyotrophic lateral sclerosis, convulsions, seizures, Huntington's disease, amnesia, or damage to the nervous system resulting from reduced oxygen supply, poison, or other toxic substances (See, e.g., U.S. Pat. No. 5,312,928).

Voltage gated calcium channels have been classified by their electrophysiological and pharmacological properties as T, L, N, P and Q types (for reviews see McCleskey *et al.* (1991) *Curr. Topics Membr.* 39:295-326; and Dunlap *et al.* (1995) *Trends. Neurosci.* 18:89-98). Because there is some overlap in the biophysical properties of the high voltage-activated channels, pharmacological profiles are useful to further distinguish them. L-type channels are sensitive to dihydropyridine agonists and antagonists. N-type channels are blocked by the peptides ω -conotoxin GVIA and ω -conotoxin MVIIA, peptide toxins from the cone shell mollusks, *Conus geographus* and *Conus magus*, respectively. P-type channels are blocked by the peptide ω -agatoxin IVA from the venom of the funnel web spider, *Agelenopsis aperta*, although some studies have suggested that ω -agatoxin IVA also blocks N-type channels (Sidach *et al.* (2000) *J. Neurosci.* 20: 7174-82). A fourth type of high voltage-activated calcium channel (Q-type) has been described, although whether the Q- and P-type channels are distinct molecular entities is controversial (Sather *et al.* (1995) *Neuron* 11:291-303; Stea *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:10576-10580; Bourinet *et al.* (1999) *Nature Neuroscience* 2:407-415).


Voltage gated calcium channels are primarily defined by the combination of different subunits: α_1 , α_2 , β , γ , and δ (see Caterall (2000) *Annu. Rev. Cell. Dev. Biol.* 16: 521-55). Ten types of α_1 subunits, four complexes, four β subunits, and two γ subunits are known (see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra; see also Klugbauer et al. (1999) *J. Neurosci.* 19: 684-691).

Based upon the combination of different subunits, calcium channels may be divided into three structurally and functionally related families: Ca_v1 , Ca_v2 , and Ca_v3 (for reviews, see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra; Ertel et al. (2000) *Neuron* 25: 533-55). L-type currents are mediated by a Ca_v1 family of α_1 subunits (see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra). Ca_v2 channels form a distinct family with less than 40% amino acid sequence identity with $Ca_v1\alpha_1$ subunits (see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra). Cloned $Ca_v2.1$ subunits conduct P- or Q-type currents that are inhibited by ω -agatoxin IVA (see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra; Sather et al. (1993) *Neuron* 11: 291-303; Stea et al. (1994) *Proc. Natl. Acad. Sci. USA* 91: 10576-80; Bourinet et al. (1999) *Nat. Neurosci.* 2: 407-15). $Ca_v2.2$ subunits conduct N-type calcium currents and have a high affinity for ω -conotoxin GVIA, ω -conotoxin MVIIA, and synthetic versions of these peptides including Ziconotide (see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra; Dubel et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:5058-62; Williams et al. (1992) *Science* 257: 389-95). Cloned $Ca_v2.3$ subunits conduct a calcium current known as R-type and are resistant to organic antagonists specific for L-type calcium currents and peptide toxins specific for N-type or P/Q-type currents (see Caterall, *Annu. Rev. Cell. Dev. Biol.*, supra; Randall et al. (1995) *J. Neurosci.* 15: 2995-3012; Soong et al. (1994) *Science* 260: 1133-36; Zhang et al. (1993) *Neuropharmacology* 32: 1075-88).


Gamma-aminobutyric acid (GABA) analogs are compounds that are derived from or based on GABA. GABA analogs are either readily available or readily synthesized using methodologies known to those of skill in the art. Exemplary GABA analogs and their salts include gabapentin and pregabalin, and any other GABA analogs as described in U.S. Pat. No. 4,024,175, U.S. Pat. No. 5,563,175, U.S. Patent No. 6,316,638, PCT Publication No. WO 93/23383, Bryans et al. (1998) *J. Med. Chem.* 41:1838-1845, and Bryans et al. (1999) *Med. Res. Rev.* 19:149-177, which are hereby incorporated by

reference. Agents useful in the practice of the invention also include those disclosed in U.S. Application No. 20020111338, cyclic amino acid compounds as disclosed in PCT Publication No. WO 99/08670, compositions disclosed in PCT Publication No. WO 99/08670, U.S. Patent No. 6,342,529, controlled release formulations as disclosed in U.S. Application No. 20020119197 and U.S. Patent No. 5,955,103, and sustained release compounds and formulations as disclosed in PCT Publication No. WO 02/28411, PCT Publication No. WO 02/28881, PCT Publication No. WO 02/28883, PCT Publication No. WO 02/32376, PCT Publication No. WO 02/42414, U.S. Application No. 20020107208, U.S. Application No. 20020151529, and U.S. Application No. 20020098999.

Gabapentin (Neurontin, or 1-(aminomethyl) cyclohexanecarboxylic acid) is an anticonvulsant drug with a high binding affinity for some calcium channel subunits, and is represented by the following structure:



Gabapentin is one of a series of compounds of formula:



in which R_1 is hydrogen or a lower alkyl radical and n is 4, 5, or 6. Although gabapentin was originally developed as a GABA-mimetic compound to treat spasticity, gabapentin has no direct GABAergic action and does not block GABA uptake or metabolism. (For review, see Rose *et al.* (2002) *Analgesia* 57:451-462). Gabapentin has been found, however, to be an effective treatment for the prevention of partial seizures in patients who are refractory to other anticonvulsant agents (Chadwick (1991) *Gabapentin*, In Pedley T A, Meldrum B S (eds.), *Recent Advances in Epilepsy*, Churchill Livingstone, New York, pp. 211-222). Gabapentin and the related drug pregabalin may interact with the $\alpha_2\delta$ subunit of calcium channels (Gee *et al.* (1996) *J. Biol. Chem.* 271: 5768-5776).


In addition to its known anticonvulsant effects, gabapentin has been shown to block the tonic phase of nociception induced by formalin and carrageenan, and exerts an inhibitory effect in neuropathic pain models of mechanical hyperalgesia and mechanical/thermal allodynia (Rose *et al.* (2002) *Analgesia* 57: 451-462). Double-blind, placebo-controlled trials have indicated that gabapentin is an effective treatment for painful symptoms associated with diabetic peripheral neuropathy, post-herpetic neuralgia, and neuropathic pain (see, e.g., Backonja *et al.* (1998) *JAMA* 280:1831-1836; Mellekers *et al.* (2001) *Clin. J. Pain* 17:284-95).

Pregabalin, (S)-(3-aminomethyl)-5-methylhexanoic acid or (S)-isobutyl GABA, is another GABA analog whose use as an anticonvulsant has been explored (Bryans *et al.* (1998) *J. Med. Chem.* 41:1838-1845). Pregabalin has been shown to possess even higher binding affinity for the $\alpha_2\delta$ subunit of calcium channels than gabapentin (Bryans *et al.* (1999) *Med. Res. Rev.* 19:149-177).



Other GABA analogs which display binding affinity to the $\alpha_2\delta$ subunit of calcium channels include, without limitation, cis-(1S,3R)-(1-(aminomethyl)-3-methylcyclohexane)acetic acid, cis-(1R,3S)-(1-(aminomethyl)-3-methylcyclohexane)acetic acid, 1 α ,3 α ,5 α -(1-(aminomethyl)-(3,5-dimethylcyclohexane)acetic acid, (9-(aminomethyl)bicyclo[3.3.1]non-9-yl)acetic acid, and (7-(aminomethyl)bicyclo[2.2.1]hept-7-yl)acetic acid (Bryans *et al.* (1998) *J. Med. Chem.* 41:1838-1845; Bryans *et al.* (1999) *Med. Res. Rev.* 19:149-177).

Fused bicyclic or tricyclic amino acid analogs of gabapentin have also been identified that are useful in the present invention. Such compounds include, for example:





1. Cyclic amino acids (illustrated below) as disclosed in PCT Publication No. WO99/21824 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;




2. Bicyclic amino acids (illustrated below) as disclosed in published U.S. Patent Application No. 60/160725, including those disclosed as having high activity as measured in a radioligand binding assay using [3H]gabapentin and the $\alpha_2\delta$ subunit derived from porcine brain tissue or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and



3. Bicyclic amino acid analogs (illustrated below) as disclosed in UK Patent Application GB 2 374 595 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.






Other agents useful in the present invention include any compound that binds to the $\alpha_2\delta$ subunit of a calcium channel. Compounds that have been identified as modulators of calcium channels include, but are not limited to those described in US Patent No. 6,316,638, US Patent No. 6,492,375, US Patent No. 6,294,533, US Patent No. 6,011,035, US Patent No. 6,387,897, US Patent No. 6,310,059, US Patent No. 6,294,533, US Patent No. 6,267,945, PCT Publication No. WO01/49670, PCT Publication No. WO01/46166, and PCT Publication No. WO01/45709. The identification of which of these compounds have a binding affinity for the $\alpha_2\delta$ subunit of calcium channels can be determined by performing $\alpha_2\delta$ binding affinity studies as described by Gee *et al.* (Gee *et al.* (1996) *J. Biol. Chem.* 271:5768-5776). The identification of still further compounds, including other GABA analogs, that have a binding affinity for the $\alpha_2\delta$ subunit of calcium channels can also be determined by performing $\alpha_2\delta$ binding affinity studies as described by Gee *et al.* (Gee *et al.* (1996) *J. Biol. Chem.* 271:5768-5776).

Acetylcholine is a chemical neurotransmitter in the nervous systems of all animals. "Cholinergic neurotransmission" refers to neurotransmission that involves acetylcholine, and has been implicated in the control of functions as diverse as locomotion, digestion, cardiac rate, "fight or flight" responses, and learning and memory (Salvatera (Feb. 2000) Acetylcholine. In *Encyclopedia of Life Sciences*. London: Nature Publishing Group, <http://www.els.net>). Receptors for acetylcholine are classified into two

general categories based on the plant alkaloids that preferentially interact with them: 1) nicotinic (nicotine binding); or 2) muscarinic (muscarine binding) (See, e.g., Salvaterra, Acetylcholine, supra).

The two general categories of acetylcholine receptors may be further divided into subclasses based upon differences in their pharmacological and electrophysiological properties. For example, nicotinic receptors are composed of a variety of subunits that are used to identify the following subclasses: 1) muscle nicotinic acetylcholine receptors; 2) neuronal nicotinic acetylcholine receptors that do not bind the snake venom α -bungarotoxin; and 3) neuronal nicotinic acetylcholine receptors that do bind the snake venom α -bungarotoxin (Dani et al. (July 1999) Nicotinic Acetylcholine Receptors in Neurons. In *Encyclopedia of Life Sciences*. London: Nature Publishing Group, <http://www.els.net>; Lindstrom (October 2001) Nicotinic Acetylcholine Receptors. In *Encyclopedia of Life Sciences*. London: Nature Publishing Group, <http://www.els.net>). By contrast, muscarinic receptors may be divided into five subclasses, labeled M_1 - M_5 , and preferentially couple with specific G-proteins (M_1 , M_3 , and M_5 with G_q ; M_2 and M_4 with G_i/G_o) (Nathanson (July 1999) Muscarinic Acetylcholine Receptors. In *Encyclopedia of Life Sciences*. London: Nature Publishing Group, <http://www.els.net>).

Other agents useful in the present invention include any anticholinergic agent, specifically, any antimuscarinic agent. Particularly useful in the methods of the present invention is oxybutynin, also known as 4-diethylaminio-2-butynyl phenylcyclohexylglycolate. It has the following structure:



Ditropan[®] (oxybutynin chloride) is the d,l racemic mixture of the above compound, which is known to exert antispasmodic effect on smooth muscle and inhibit the muscarinic action of acetylcholine on smooth muscle. Metabolites and isomers of oxybutynin have also been shown to have activity useful according to the present invention. Examples include, but are not limited to N-desethyl-oxybutynin and S-oxybutynin.

Additional compounds that have been identified as antimuscarinic agents and are useful in the present invention include, but are not limited to:

- a. Darifenacin (Daryon[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Solifenacin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. YM-905 (solifenacin succinate) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. Solifenacin monohydrochloride or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Tolterodine (Detrol[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. Propiverine (Detrunorm[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. Propantheline bromide (Pro-Banthine[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. Hyoscyamine sulfate (Levsin[®], Cystospaz[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Dicyclomine hydrochloride (Bentyl[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Flavoxate hydrochloride (Urispas[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- k. d,l (racemic) 4- diethylamino-2-butynyl phenylcyclohexylglycolate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. (R)-N,N-diisopropyl-3-(2-hydroxy-5-methylphenyl)-3-phenylpropanamine L-hydrogen tartrate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. (+)-(1S,3'R)-quinuclidin-3'-yl 1-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate monosuccinate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. alpha(+)-4-(Dimethylamino)-3-methyl-1,2-diphenyl-2-butanol propionate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. 1-methyl-4-piperidyl diphenylpropoxyacetate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. 3'-hydroxyspiro[1'H,5'H-nortropane-8,1'-pyrrolidinium benzilate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. 4-amino-piperidine containing compounds as disclosed in Diouf *et al.* (2002) *Bioorg. Med. Chem. Lett.* 12: 2535-9 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- r. Pirenzepine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. Methoctramine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- t. 4-diphenylacetoxy-N-methyl piperidine methiodide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- u. Tropicamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- v. (2R)-N-[1-(6-aminopyridin-2-ylmethyl)piperidin-4-yl]-2-[(1R)-3,3-difluorocyclopentyl]-2-hydroxy-2-phenylacetamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. PNU-200577 ((R)-N, N-diisopropyl-3-(2-hydroxy-5-hydroxymethylphenyl)-3-phenylpropanamine) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. KRP-197(4-(2-methylimidazolyl)-2,2-diphenylbutyramide) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof
- y. Fesoterodine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- z. SPM 7605 (the active metabolite of Fesoterodine), or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof

The identification of further compounds that have antimuscarinic activity and would therefore be useful in the present invention can be determined by performing muscarinic receptor binding specificity studies as described by Nilvebrant (2002) *Pharmacol. Toxicol.* 90: 260-7 or cystometry studies as described by Modiri *et al.* (2002) *Urology* 59: 963-8.

Adrenergic receptors are cell-surface receptors for two major catecholamine hormones and neurotransmitters: noradrenaline and adrenaline. (Malbon *et al.* (Feb. 2000) Adrenergic Receptors. In *Encyclopedia of Life Sciences*. London: Nature Publishing Group, <http://www.els.net>). Adrenergic receptors have been implicated in

critical physiological processes, including blood pressure control, myocardial and smooth muscle contractility, pulmonary function, metabolism, and central nervous system activity (See, e.g., Malbon *et al.*, Adrenergic Receptors, *supra*). Two classes of adrenergic receptors have been identified, α and β , that may be further subdivided into three major families ($\alpha 1$, $\alpha 2$, and β), each with at least three subtypes ($\alpha 1A$, B, and D; $\alpha 2A$, B, and C; and $\beta 1$, $\beta 2$, and $\beta 3$) based upon their binding characteristics to different agonists and molecular cloning techniques. (See, e.g., Malbon *et al.*, Adrenergic Receptors, *supra*).

Other agents useful in the present invention include any $\beta 3$ adrenergic agonist agent. Compounds that have been identified as $\beta 3$ adrenergic agonist agents and are useful in the present invention include, but are not limited to:

- a. TT-138 and phenylethanolamine compounds as disclosed in US Patent No. 6,069,176, PCT Publication No. WO 97/15549 and available from Mitsubishi Pharma Corp. or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- b. FR-149174 and propanolamine derivatives as disclosed in US Patent Nos. 6,495,546 and 6,391,915 and available from Fujisawa Pharmaceutical Co. or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- c. KUC-7483, available from Kissei Pharmaceutical Co. or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- d. 4'-hydroxynorephedrine derivatives such as 2-2-chloro-4-(2-(1S,2R)-2-hydroxy-2-(4-hydroxyphenyl)-1-methylethylamino)ethyl)phenoxy acetic acid as disclosed in Tanaka *et al.* (2003) *J. Med. Chem.* 46: 105-12 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- e. 2-amino-1-phenylethanol compounds, such as BRL35135 ((R*R*)-(.-.-)-[4-[2-[2-(3-chlorophenyl)-2-

- ydroxyethylamino]propyl]phenox y]acetic acid methyl ester hydrobromide salt as disclosed in Japanese Patent Publication No. 26744 of 1988 and European Patent Publication No. 23385), and SR58611A ((RS)-N-(7-ethoxycarbonylmethoxy-1,2,3,4-tetrahydronaphth-2-yl)-2-(3-chlor ophenyl)-2-hydroxyethanamine hydrochloride as disclosed in Japanese Laid-open Patent Publication No. 66152 of 1989 and European Laid-open Patent Publication No. 255415) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- f. GS 332 (Sodium (2R)-[3-[3-[2-(3 Chlorophenyl)-2-hydroxyethylamino]cyclohexyl]phenoxy]acetate) as disclosed in Iizuka *et al.* (1998) *J. Smooth Muscle Res.* 34: 139-49 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- g. BRL-37,344 (4-[-(2-hydroxy-(3-chlorophenyl) ethyl)-amino]propyl]phenoxyacetate) as disclosed in Tsujii *et al.* (1998) *Physiol. Behav.* 63: 723-8 and available from Glaxosmithkline or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- h. BRL-26830A as disclosed in Takahashi *et al.* (1992) *Jpn Circ. J.* 56: 936-42 and available from Glaxosmithkline or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- i. CGP 12177 (4-[3-t-butylamino-2-hydroxypropoxy]benzimidazol-2- one) (a $\beta 1/\beta 2$ adrenergic antagonist reported to act as an agonist for the $\beta 3$ adrenergic receptor) as described in Tavernier *et al.* (1992) *J. Pharmacol. Exp. Ther.* 263: 1083-90 and available from Ciba-Geigy or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;

- j. CL 316243 (R,R-5-[2-[[2-(3-chlorophenyl)-2-hydroxyethyl]amino]propyl]-1,3- benzodioxole-2,2-dicarboxylate) as disclosed in Berlan *et al.* (1994) *J. Pharmacol. Exp. Ther.* 268: 1444-51 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- k. Compounds having β_3 adrenergic agonist activity as disclosed in US Patent Application 20030018061 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- l. ICI 215,001 HCl ((S)-4-[2-Hydroxy-3-phenoxypropylaminoethoxy]phenoxyacetic acid hydrochloride) as disclosed in Howe (1993) *Drugs Future* 18: 529 and available from AstraZeneca/ICI Labs or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- m. ZD 7114 HCl (ICI D7114; (S)-4-[2-Hydroxy-3-phenoxypropylaminoethoxy]-N-(2-methoxyethyl)phenoxyacetamide HCl) as disclosed in Howe (1993) *Drugs Future* 18: 529 and available from AstraZeneca/ICI Labs or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- n. Pindolol (1-(1H-Indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol) as disclosed in Blin *et al* (1994) *Mol.Pharmacol.* 44: 1094 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- o. (S)-(-)-Pindolol ((S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol) as disclosed in Walter *et al* (1984) *Naunyn-Schmied.Arch.Pharmacol.* 327: 159 and Kalkman (1989) *Eur.J.Pharmacol.* 173: 121 or acids, salts,

- esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- p. SR 59230A HCl (1-(2-Ethylphenoxy)-3-[[[(1S)-1,2,3,4-tetrahydro-1-naphthalenyl]amino]-(2S)-2-propanol hydrochloride) as disclosed in Manara *et al.* (1995) *Pharmacol. Comm.* 6: 253 and Manara *et al.* (1996) *Br. J. Pharmacol.* 117: 435 and available from Sanofi-Midy or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- q. SR 58611 (N[2s]7-carb-ethoxymethoxy-1,2,3,4-tetrahydronaphthyl)-(2r)-2-hydroxy-2(3-chlorophenyl) ethamine hydrochloride) as disclosed in Gauthier *et al.* (1999) *J. Pharmacol. Exp. Ther.* 290: 687-693 and available from Sanofi Research or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof; and
- r. YM178 available from Yamanouchi Pharmaceutical Co. or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof.

The identification of further compounds that have β_3 adrenergic agonist activity and would therefore be useful in the present invention can be determined by performing radioligand binding assays and/or contractility studies as described by Zilberfarb *et al.* (1997) *J. Cell Sci.* 110: 801-807; Takeda *et al.* (1999) *J. Pharmacol. Exp. Ther.* 288: 1367-1373; and Gauthier *et al.* (1999) *J. Pharmacol. Exp. Ther.* 290: 687-693.

Spasmolytics are compounds that relieve or prevent muscle spasms. Compounds that have been identified as spasmolytic agents and that are useful in the present invention include, but are not limited to:

- a. α - α -diphenylacetic acid-4-(N-methyl-piperidyl) esters as disclosed in US Patent No. 5,897,875 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;

- b. Human and porcine spasmolytic polypeptides in glycosylated form and variants thereof as disclosed in US Patent No. 5,783,416 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- c. Dioxazocine derivatives as disclosed in US Patent No. 4,965,259 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- d. Quaternary 6,11-dihydro-dibenzo-[b,e]-thiepine-11-N-alkylnorscopine ethers as disclosed in US Patent No. 4,608,377 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- e. Quaternary salts of dibenzo[1,4]diazepinones, pyrido-[1,4]benzodiazepinones, pyrido[1,5]benzodiazepinones as disclosed in US Patent No. 4,594,190 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- f. Endo-8,8-dialkyl-8-azoniabicyclo (3.2.1) octane-6,7-exo-epoxy-3-alkyl-carboxylate salts as disclosed in US Patent No. 4,558,054 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- g. Pancreatic spasmolytic polypeptides as disclosed in US Patent No. 4,370,317 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- h. Triazinones as disclosed in US Patent No. 4,203,983 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- i. 2-(4-Biphenyl)-N-(2-diethylamino alkyl)propionamide as disclosed in US Patent No. 4,185,124 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;

- j. Piperazino-pyrimidines as disclosed in US Patent No. 4,166,852 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- k. Aralkylamino carboxylic acids as disclosed in US Patent No. 4,163,060 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- l. Aralkylamino sulfones as disclosed in US Patent No. 4,034,103 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- m. Smooth muscle spasmolytic agents as disclosed in US Patent No. 6,207,852 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof; and
- n. Papaverine or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof.

The identification of further compounds that have spasmolytic activity and would therefore be useful in the present invention can be determined by performing contractility studies as described in US Patent No. 6,207,852; Noronha-Blob *et al.* (1991) *J. Pharmacol. Exp. Ther.* 256: 562-567; and/or Kachur *et al.* (1988) *J. Pharmacol. Exp. Ther.* 247: 867-872.

Tachykinins (TKs) are a family of structurally related peptides that include substance P, neurokinin A (NKA) and neurokinin B (NKB). Neurons are the major source of TKs in the periphery. An important general effect of TKs is neuronal stimulation, but other effects include endothelium-dependent vasodilation, plasma protein extravasation, mast cell recruitment and degranulation and stimulation of inflammatory cells (See Maggi, C. A. (1991) *Gen. Pharmacol.*, 22: 1-24).

Substance P activates the neurokinin receptor subtype referred to as NK₁. Substance P is an undecapeptide that is present in sensory nerve terminals. Substance P is known to have multiple actions that produce inflammation and pain in the periphery after C-fiber activation, including vasodilation, plasma extravasation and degranulation of mast cells (Levine, J. D. *et al.* (1993) *J. Neurosci.* 13: 2273). Neurokinin A is a peptide which is colocalized in sensory neurons with substance P and which also

promotes inflammation and pain. Neurokinin A activates the specific neurokinin receptor referred to as NK₂ (Edmonds-Alt, S., *et. al.* (1992) *Life Sci.* 50: PL101). Suitable neurokinin receptor antagonists for use in the present invention that act on the NK₁ receptor include, but are not limited to: 1-imino-2-(2-methoxy-phenyl)-ethyl-7,7-diphenyl-4-perhydroisoindolone(3aR, 7aR) ("RP 67580"); 2S,3S-cis-3-(2-methoxybenzylamino)-2-benzhydrylquinuclidine ("CP 96,345"); and (aR,9R)-7-[3,5-bis(trifluoromethyl)benzyl]-8,9,10, 11-tetrahydro-9-methyl-5-(4-methylphenyl)-7H-[1,4]diazocino[2,1-g] [1,7]naphthyridine-6,13-dione)("TAK-637"). Suitable neurokinin receptor antagonists for use in the present invention that act on the NK₂ receptor include but are not limited to: ((S)-N-methyl-N-4-(4-acetylamino-4-phenylpiperidino)-2-(3,4-dichlorophenyl)butylbenzamide ("SR 48968"); Met-Asp-Trp-Phe-Dap-Leu ("MEN 10,627"); and cyc(Gln-Trp-Phe-Gly-Leu-Met) ("L 659,877"). The identification of further compounds that have neurokinin receptor antagonist activity and would therefore be useful in the present invention can be determined by performing binding assay studies as described in Hopkins *et al.* (1991) *Biochem. Biophys. Res. Comm.* 180: 1110-1117; and Aharony *et al.* (1994) *Mol. Pharmacol.* 45: 9-19.

Bradykinin receptors generally are divided into bradykinin₁ (B₁) and bradykinin₂ (B₂) subtypes. Studies have shown that acute peripheral pain and inflammation produced by bradykinin are mediated by the B₂ subtype whereas bradykinin-induced pain in the setting of chronic inflammation is mediated via the B₁ subtype (Perkins, M. N., *et. al.* (1993) *Pain* 53: 191-97); Dray, A., *et. al.* (1993) *Trends Neurosci.* 16: 99-104). Suitable bradykinin receptor antagonists for use in the present invention that act on the B₁ receptor include but are not limited to: des-arg¹⁰HOE 140 (available from Hoechst Pharmaceuticals) and des-Arg⁹bradykinin (DABK). Suitable bradykinin receptor antagonists for use in the present invention that act on the B₂ receptor include but are not limited to: D-Phe⁷-BK; D-Arg-(Hyp³-Thi^{5,8}-D-Phe⁷)-BK ("NPC 349"); D-Arg-(Hyp³-D-Phe⁷)-BK ("NPC 567"); D-Arg-(Hyp³-Thi⁵-D-Tic⁷-Oic⁸)-BK ("HOE 140"); H-DArg-Arg-Pro-Hyp-Gly-Thi-c(Dab-DTic-Oic-Arg)c(7gamma-10alpha)("MEN11270"); H-DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg-OH("Icatibant"); (E)-3-(6-acetamido-3-

pyridyl)-N-[N-[2, 4-dichloro-3-[(2-methyl-8-quinolinyloxy)methyl]phenyl]-N-methylaminocarbonylmethyl]acrylamide ("FR173567"); and WIN 64338.

These compounds are more fully described in Perkins, M. N., *et al.*, Pain, *supra*; Dray, A., *et al.*, *Trends Neurosci.*, *supra*; and Meini *et al.* (2000) *Eur. J. Pharmacol.* 388: 177-82. The identification of further compounds that have bradykinin receptor antagonist activity and would therefore be useful in the present invention can be determined by performing binding assay studies as described in Manning *et al.* (1986) *J. Pharmacol. Exp. Ther.* 237: 504 and US Patent No. 5,686,565.

Nitric oxide donors may be included in the present invention particularly for their anti-spasm activity. Nitric oxide (NO) plays a critical role as a molecular mediator of many physiological processes, including vasodilation and regulation of normal vascular tone. The action of NO is implicated in intrinsic local vasodilation mechanisms. NO is the smallest biologically active molecule known and is the mediator of an extraordinary range of physiological processes (Nathan (1994) *Cell* 78: 915-918; Thomas (1997) *Neurosurg. Focus* 3: Article 3). NO is also a known physiologic antagonist of endothelin-1, which is the most potent known mammalian vasoconstrictor, having at least ten times the vasoconstrictor potency of angiotensin II (Yanagisawa *et al.* (1988) *Nature* 332: 411-415; Kasuya *et al.* (1993) *J. Neurosurg.* 79: 892-898; Kobayashi *et al.*, (1991) *Neurosurgery* 28: 673-679). The biological half-life of NO is extremely short (Morris *et al.* (1994) *Am. J. Physiol.* 266: E829-E839; Nathan (1994) *Cell* 78: 915-918). NO accounts entirely for the biological effects of endothelium-derived relaxing factor (EDRF) and is an extremely potent vasodilator that is believed to work through the action of cGMP-dependent protein kinases to effect vasodilation (Henry *et al.* (1993) *FASEB J.* 7: 1124-1134; Nathan (1992) *FASEB J.* 6: 3051-3064; Palmer *et al.*, (1987) *Nature* 327: 524-526; Snyder *et al.* (1992) *Scientific American* 266: 68-77).

Within endothelial cells, an enzyme known as NO synthase (NOS) catalyzes the conversion of L-arginine to NO which acts as a diffusible second messenger and mediates responses in adjacent smooth muscle cells. NO is continuously formed and released by the vascular endothelium under basal conditions which inhibits contractions and controls basal coronary tone and is produced in the endothelium in response to various agonists (such as acetylcholine) and other endothelium dependent vasodilators.

Thus, regulation of NOS activity and the resultant levels of NO are key molecular targets controlling vascular tone (Muramatsu *et. al.* (1994) *Coron. Artery Dis.* 5: 815-820).

Other agents useful in the present invention include any nitric oxide donor agent. Suitable nitric oxide donors for the practice of the present invention include but are not limited to:

- a. Nitroglycerin or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- b. Sodium nitroprusside or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- c. FK 409 (NOR-3) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- d. FR 144420 (NOR-4) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- e. 3-morpholiniosydnonimine or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- f. Linsidomine chlorohydrate ("SIN-1") or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- g. S-nitroso-N-acetylpenicillamine ("SNAP") or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- h. AZD3582 (CINOD lead compound, available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- i. NCX 4016 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- j. NCX 701 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;

- k. NCX 1022 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- l. HCT 1026 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- m. NCX 1015 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- n. NCX 950 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- o. NCX 1000 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- p. NCX 1020 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- q. AZD 4717 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- r. NCX 1510/NCX 1512 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- s. NCX 2216 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- t. NCX 4040 (available from NicOx S.A.) or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;

- u. Nitric oxide donors as disclosed in U.S. Patent No. 5,155,137 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- v. Nitric oxide donors as disclosed in U.S. Patent No. 5,366,997 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- w. Nitric oxide donors as disclosed in U.S. Patent No. 5,405,919 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- x. Nitric oxide donors as disclosed in U.S. Patent No. 5,650,442 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- y. Nitric oxide donors as disclosed in U.S. Patent No. 5,700,830 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- z. Nitric oxide donors as disclosed in U.S. Patent No. 5,632,981 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- aa. Nitric oxide donors as disclosed in U.S. Patent No. 6,290,981 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- bb. Nitric oxide donors as disclosed in U.S. Patent No. 5,691,423 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- cc. Nitric oxide donors as disclosed in U.S. Patent No. 5,721,365 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;
- dd. Nitric oxide donors as disclosed in U.S. Patent No. 5,714,511 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof;

- ee. Nitric oxide donors as disclosed in U.S. Patent No. 6,511,911 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof; and
- ff. Nitric oxide donors as disclosed in U.S. Patent No. 5,814,666 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof.

The identification of further compounds that have nitric oxide donor activity and would therefore be useful in the present invention can be determined by release profile and/or induced vasospasm studies as described in US Patent Nos. 6,451,337 and 6,358,536, as well as Moon (2002) *IBJU Int.* 89: 942-9 and Fathian-Sabet *et al.* (2001) *J. Urol.* 165: 1724-9.

Enantiomers and Diastereomers


Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane-polarized light. In describing an optically active compound the prefixes R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes D and L, or (+) or (-), designate the sign of rotation of plane-polarized light by the compound, with L or (-) meaning that the compound is levorotatory. In contrast, a compound prefixed with D or (+) is dextrorotatory. There is no correlation between nomenclature for the absolute stereochemistry and for the rotation of an enantiomer. Thus, D-lactic acid is the same as (-)-lactic acid, and L-lactic acid is the same as (+)-lactic acid. For a given chemical structure, each of a pair of enantiomers are identical except that they are non-superimposable mirror images of one another. A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric, or racemic, mixture.

Stereochemical purity is important in the pharmaceutical field, where many of the most often prescribed drugs exhibit chirality. For example, the L-enantiomer of the beta-adrenergic blocking agent, propranolol, is known to be 100 times more potent than its D-enantiomer. Additionally, optical purity is important in the pharmaceutical drug field because certain isomers have been found to impart a deleterious effect, rather than an advantageous or inert effect. For example, it is believed that the D-enantiomer of

thalidomide is a safe and effective sedative when prescribed for the control of morning sickness during pregnancy, whereas its corresponding L-enantiomer is believed to be a potent teratogen.

When two chiral centers exist in one molecule, there are four possible stereoisomers: (R,R), (S,S), (R,S), and (S,R). Of these, (R,R) and (S,S) are an example of a pair of enantiomers (mirror images of each other), which typically share chemical properties and melting points just like any other enantiomeric pair. The mirror images of (R,R) and (S,S) are not, however, superimposable on (R,S) and (S,R). This relationship is called diastereoisomeric, and the (S,S) molecule is a diastereoisomer of the (R,S) molecule, whereas the (R,R) molecule is a diastereoisomer of the (S,R) molecule.

An example of a compound with two chiral centers is the antimuscarinic solifenacin. Solifenacin is described in U.S. Patent No. 6,174,896 and is represented by the following chemical formula:



Because solifenacin has two chiral centers, diastereomers as well as enantiomers exist for this molecule (see U.S. Patent No. 6,174,896). Solifenacin succinate (development number YM-905) is a salt form of solifenacin that is co-promoted as Vesicare[®] by Yamanouchi Pharmaceutical Co., Ltd. (through Yamanouchi Pharma America) and GlaxoSmithKline as an investigational muscarinic antagonist. Solifenacin was discovered and developed by Yamanouchi, and a New Drug Application was submitted to the U.S. Food and Drug Administration by YPA in December 2002 for solifenacin succinate. A market authorization application for Vesicare[®] was submitted in Europe in January 2003, and Yamanouchi has initiated Phase III clinical trials for Vesicare[®] in Japan. Other salt forms of solifenacin have also been specifically described by Yamanouchi, including solifenacine monohydrochloride (development number YM-53705).

For use in the present invention, any diastereomer or enantiomer of an active agent, as disclosed herein, can be administered to treat functional bowel disorders in patients in need of such treatment.

Formulations

Formulations of the present invention may include, but are not limited to, continuous, as needed, short-term, rapid-offset, controlled release, sustained release, delayed release, and pulsatile release formulations.

Compositions of the invention comprise $\alpha_2\delta$ subunit calcium channel modulators in combination with one or more compounds with smooth muscle modulatory effects, including antimuscarinics (particularly those that do not have an amine embedded in an 8-azabicyclo[3.2.1]octan-3-ol skeleton), β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors. The compositions are administered in therapeutically effective amounts to a patient in need thereof for treating functional bowel disorders, particularly IBS. It is recognized that the compositions may be administered by any means of administration as long as an effective amount for the treatment of functional bowel disorders is delivered.

Any of the active agents may be administered in the form of a salt, ester, amide, prodrug, active metabolite, derivative, or the like, provided that the salt, ester, amide, prodrug or derivative is suitable pharmacologically, i.e., effective in the present method. Salts, esters, amides, prodrugs and other derivatives of the active agents may be prepared using standard procedures known to those skilled in the art of synthetic organic chemistry and described, for example, by J. March, *Advanced Organic Chemistry: Reactions, Mechanisms and Structure*, 4th Ed. (New York: Wiley-Interscience, 1992). For example, acid addition salts are prepared from the free base using conventional methodology, and involves reaction with a suitable acid. Suitable acids for preparing acid addition salts include both organic acids, e.g., acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like, as well as inorganic acids, e.g., hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid,

phosphoric acid, and the like. An acid addition salt may be reconverted to the free base by treatment with a suitable base. Particularly preferred acid addition salts of the active agents herein are salts prepared with organic acids. Conversely, preparation of basic salts of acid moieties which may be present on an active agent are prepared in a similar manner using a pharmaceutically acceptable base such as sodium hydroxide, potassium hydroxide, ammonium hydroxide, calcium hydroxide, trimethylamine, or the like.

Preparation of esters involves functionalization of hydroxyl and/or carboxyl groups that may be present within the molecular structure of the drug. The esters are typically acyl-substituted derivatives of free alcohol groups, i.e., moieties that are derived from carboxylic acids of the formula RCOOH where R is alkyl, and preferably is lower alkyl. Esters can be reconverted to the free acids, if desired, by using conventional hydrogenolysis or hydrolysis procedures. Amides and prodrugs may also be prepared using techniques known to those skilled in the art or described in the pertinent literature. For example, amides may be prepared from esters, using suitable amine reactants, or they may be prepared from an anhydride or an acid chloride by reaction with ammonia or a lower alkyl amine. Prodrugs are typically prepared by covalent attachment of a moiety, which results in a compound that is therapeutically inactive until modified by an individual's metabolic system.

One set of formulations for gabapentin are those marketed by Pfizer Inc. under the brand name Neurontin®. Neurontin® Capsules, Neurontin® Tablets, and Neurontin® Oral Solution are supplied either as imprinted hard shell capsules containing 100 mg, 300 mg, and 400 mg of gabapentin, elliptical film-coated tablets containing 600 mg and 800 mg of gabapentin or an oral solution containing 250 mg/5 mL of gabapentin. The inactive ingredients for the capsules are lactose, cornstarch, and talc. The 100 mg capsule shell contains gelatin and titanium dioxide. The 300 mg capsule shell contains gelatin, titanium dioxide, and yellow iron oxide. The 400 mg capsule shell contains gelatin, red iron oxide, titanium dioxide, and yellow iron oxide. The inactive ingredients for the tablets are poloxamer 407, copolyvidonum, cornstarch, magnesium stearate, hydroxypropyl cellulose, talc, candelilla wax and purified water. The inactive ingredients for the oral solution are glycerin, xylitol, purified water and artificial cool strawberry anise flavor. In addition to these formulations, gabapentin and formulations are generally

described in the following patents: US 6,683,112; US 6,645,528; US 6,627,211; US 6,569,463; US 6,544,998; US 6,531,509; 6,495,669; US 6,465,012; US 6,346,270; US 6,294,198; US 6,294,192; US 6,207,685; US 6,127,418; US 6,024,977; US 6,020,370; US 5,906,832; US 5,876,750; and US 4,960,931.

One set of formulations for oxybutynin are those marketed by Ortho-McNeil Pharmaceuticals, Inc. under the brand name Ditropan[®]. Ditropan[®] tablets are supplied containing 5 mg/tablets of the active ingredient, oxybutynin chloride, and the inactive ingredients anhydrous lactose, microcrystalline cellulose, calcium stearate, and FD&C blue #1 lake. Ditropan[®] syrup is supplied as 5mg/5mL of the active ingredient, oxybutynin chloride, and the inactive ingredients citric acid, FD&C green #3, flavor, glycerin, methylparaben, sodium citrate, sorbitol, sucrose, and water. Ditropan XL[®] is an extended release tablet form of Ditropan[®] supplied containing either 5 mg (pale yellow color) of oxybutynin chloride, 10 mg (pink color) of oxybutynin chloride, or 15 mg (gray color) of oxybutynin chloride. Inactive ingredients are cellulose acetate, hydroxypropyl methylcellulose, lactose, magnesium stearate, polyethylene glycol, polyethylene oxide, synthetic iron oxides, titanium dioxide, polysorbate 80, sodium chloride, and butylated hydroxytoluene.

Oxybutynin is also supplied by Watson Pharmaceuticals under the brand name Oxytrol[®] (oxybutynin transdermal system). Oxytrol[®] is a transdermal patch designed to deliver oxybutynin continuously and consistently over a 3 to 4 day interval. It is supplied as a 39 cm² patch containing 36 mg of oxybutynin, which is designed to deliver 3.9 mg/day. The patch is worn continuously, and a new patch is applied every 3 to 4 days.

A formulation useful in the present invention comprises a combination of gabapentin and oxybutynin chloride. The combination can be supplied in various pharmaceutical composition and dosage forms as described herein. One formulation for supplying the combination is in a tablet formulation. Additional formulations for the combination of the present invention, such as capsules, syrups, etc. are also envisioned for delivery of the combination, and any description of tablet formulations is in no way meant to be limiting of possible delivery modes for the combination of the present invention.

Tablet formulations useful for supplying the gabapentin/oxybutynin combination useful in the present invention can comprise, in addition to the active ingredients in combination, functional excipients. Such excipients as are useful for preparing pharmaceutical compositions in a tablet formulation are known in the art and include compounds known to be useful as fillers, binders, lubricants, disintegrants, diluents, coatings, plastizers, glidants, compression aids, stabilizers, sweeteners, solubilizers, and other excipients that would be known to one of skill in the pharmaceutical arts.

The active ingredients of the combination useful in the present invention (gabapentin and oxybutynin) can be combined, particularly in tablet form, according to ratios provided herein. The relative ratio of the active ingredients of the combination for use in the present invention is about 2.5:200 to 2.5:800, oxybutynin and gabapentin respectively. Generally, the ratio of oxybutynin to gabapentin in the combination is about 2.5:200. Alternately, the ratio of oxybutynin to gabapentin in the combination is about 2.5:400. Alternately, the ratio of oxybutynin to gabapentin in the combination is about 2.5:600. Examples of formulations for preparing tablets comprising gabapentin and oxybutynin in combination suitable for use in the present invention are provided below in Tables 1 and 2.

Table 1	
Ingredient	Weight per Unit
Gabapentin	200.0
Oxybutynin chloride	2.50
Lactose, monohydrate	85.50
Purified water	130.0
Providone	24.00
Microcrystalline cellulose	80.00
Crospovidone	4.00
Magnesium stearate	4.00
Total	400.0

Table 2	
Ingredient	Weight per Unit
Gabapentin	200.0
Oxybutynin chloride	2.50
Lactose, monohydrate	89.50
Purified water	235.0
Hydroxypropylmethylcellulose	20.00
Microcrystalline cellulose	80.00
Crospovidone	4.00
Magnesium stearate	4.00
Total	400.0

Tablets according to the above formulations can be prepared according to a number of possible methods. One method used in preparing a tablet comprising a formulation as provided above includes the following steps:

- (1) sift ingredients through 20-mesh screen, transfer to granulator with impeller and chopper, and mix for five minutes;
- (2) wet granulate mixed ingredients with a binder solution (such as povidone or methocel);
- (3) transfer wet granules to fluid bed dryer and dry until %LOD values are within a 1-2.5% range;
- (4) mill dried granules;
- (5) lubricate milled granules (such as with magnesium stearate) in blender;
- (6) compress into tablets.

Other derivatives and analogs of the active agents may be prepared using standard techniques known to those skilled in the art of synthetic organic chemistry, or may be deduced by reference to the pertinent literature. In addition, chiral active agents may be in isomerically pure form, or they may be administered as a racemic mixture of isomers.

Pharmaceutical Compositions and Dosage Forms

Suitable compositions and dosage forms include tablets, capsules, caplets, pills, gel caps, troches, dispersions, suspensions, solutions, syrups, transdermal patches, gels, powders, magmas, lozenges, creams, pastes, plasters, lotions, discs, suppositories, liquid

sprays for nasal or oral administration, dry powder or aerosolized formulations for inhalation, compositions and formulations for intravesical administration and the like. Further, those of ordinary skill in the art can readily deduce that suitable formulations involving these compositions and dosage forms, including those formulations as described elsewhere herein.

Oral Dosage Forms

Oral dosage forms include tablets, capsules, caplets, solutions, suspensions and/or syrups, and may also comprise a plurality of granules, beads, powders or pellets that may or may not be encapsulated. Such dosage forms are prepared using conventional methods known to those in the field of pharmaceutical formulation and described in the pertinent texts, e.g., in Remington: The Science and Practice of Pharmacy, supra). Tablets and capsules represent the most convenient oral dosage forms, in which case solid pharmaceutical carriers are employed.

Tablets may be manufactured using standard tablet processing procedures and equipment. One method for forming tablets is by direct compression of a powdered, crystalline or granular composition containing the active agent(s), alone or in combination with one or more carriers, additives, or the like. As an alternative to direct compression, tablets can be prepared using wet-granulation or dry-granulation processes. Tablets may also be molded rather than compressed, starting with a moist or otherwise tractable material; however, compression and granulation techniques are preferred.

In addition to the active agent(s), then, tablets prepared for oral administration using the method of the invention will generally contain other materials such as binders, diluents, lubricants, disintegrants, fillers, stabilizers, surfactants, preservatives, coloring agents, flavoring agents and the like. Binders are used to impart cohesive qualities to a tablet, and thus ensure that the tablet remains intact after compression. Suitable binder materials include, but are not limited to, starch (including corn starch and pregelatinized starch), gelatin, sugars (including sucrose, glucose, dextrose and lactose), polyethylene glycol, propylene glycol, waxes, and natural and synthetic gums, e.g., acacia sodium alginate, polyvinylpyrrolidone, cellulosic polymers (including hydroxypropyl cellulose, hydroxypropyl methylcellulose, methyl cellulose, ethyl cellulose, hydroxyethyl cellulose,

and the like), and Veegum. Diluents are typically necessary to increase bulk so that a practical size tablet is ultimately provided. Suitable diluents include dicalcium phosphate, calcium sulfate, lactose, cellulose, kaolin, mannitol, sodium chloride, dry starch and powdered sugar. Lubricants are used to facilitate tablet manufacture; examples of suitable lubricants include, for example, vegetable oils such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil, and oil of theobroma, glycerin, magnesium stearate, calcium stearate, and stearic acid. Stearates, if present, preferably represent at no more than approximately 2 wt. % of the drug-containing core. Disintegrants are used to facilitate disintegration of the tablet, and are generally starches, clays, celluloses, algin, gums or crosslinked polymers. Fillers include, for example, materials such as silicon dioxide, titanium dioxide, alumina, talc, kaolin, powdered cellulose and microcrystalline cellulose, as well as soluble materials such as mannitol, urea, sucrose, lactose, dextrose, sodium chloride and sorbitol. Stabilizers are used to inhibit or retard drug decomposition reactions that include, by way of example, oxidative reactions. Surfactants may be anionic, cationic, amphoteric or nonionic surface active agents.

The dosage form may also be a capsule, in which case the active agent-containing composition may be encapsulated in the form of a liquid or solid (including particulates such as granules, beads, powders or pellets). Suitable capsules may be either hard or soft, and are generally made of gelatin, starch, or a cellulosic material, with gelatin capsules preferred. Two-piece hard gelatin capsules are preferably sealed, such as with gelatin bands or the like. (See, for e.g., Remington: The Science and Practice of Pharmacy, supra), which describes materials and methods for preparing encapsulated pharmaceuticals. If the active agent-containing composition is present within the capsule in liquid form, a liquid carrier is necessary to dissolve the active agent(s). The carrier must be compatible with the capsule material and all components of the pharmaceutical composition, and must be suitable for ingestion.

Solid dosage forms, whether tablets, capsules, caplets, or particulates, may, if desired, be coated so as to provide for delayed release. Dosage forms with delayed release coatings may be manufactured using standard coating procedures and equipment. Such procedures are known to those skilled in the art and described in the pertinent texts (See, for e.g., Remington: The Science and Practice of Pharmacy, supra). Generally,

after preparation of the solid dosage form, a delayed release coating composition is applied using a coating pan, an airless spray technique, fluidized bed coating equipment, or the like. Delayed release coating compositions comprise a polymeric material, e.g., cellulose butyrate phthalate, cellulose hydrogen phthalate, cellulose propionate phthalate, polyvinyl acetate phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate, dioxypopyl methylcellulose succinate, carboxymethyl ethylcellulose, hydroxypropyl methylcellulose acetate succinate, polymers and copolymers formed from acrylic acid, methacrylic acid, and/or esters thereof.

Sustained release dosage forms provide for drug release over an extended time period, and may or may not be delayed release. Generally, as will be appreciated by those of ordinary skill in the art, sustained release dosage forms are formulated by dispersing a drug within a matrix of a gradually bioerodible (hydrolyzable) material such as an insoluble plastic, a hydrophilic polymer, or a fatty compound, or by coating a solid, drug-containing dosage form with such a material. Insoluble plastic matrices may be comprised of, for example, polyvinyl chloride or polyethylene. Hydrophilic polymers useful for providing a sustained release coating or matrix cellulosic polymers include, without limitation: cellulosic polymers such as hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropylmethyl cellulose phthalate, hydroxypropylcellulose phthalate, cellulose hexahydrophthalate, cellulose acetate hexahydrophthalate, and carboxymethylcellulose sodium; acrylic acid polymers and copolymers, preferably formed from acrylic acid, methacrylic acid, acrylic acid alkyl esters, methacrylic acid alkyl esters, and the like, e.g. copolymers of acrylic acid, methacrylic acid, methyl acrylate, ethyl acrylate, methyl methacrylate and/or ethyl methacrylate, with a terpolymer of ethyl acrylate, methyl methacrylate and trimethylammonioethyl methacrylate chloride (sold under the tradename Eudragit RS) preferred; vinyl polymers and copolymers such as polyvinyl pyrrolidone, polyvinyl acetate, polyvinylacetate phthalate, vinylacetate crotonic acid copolymer, and ethylene-vinyl acetate copolymers; zein; and shellac, ammoniated shellac, shellac-acetyl alcohol, and shellac n-butyl stearate. Fatty compounds for use as a sustained release matrix

material include, but are not limited to, waxes generally (e.g., carnauba wax) and glyceryl tristearate.

Transmucosal Compositions and Dosage Forms

Although the present compositions may be administered orally, other modes of administration are suitable as well. For example, transmucosal administration may be advantageously employed. Transmucosal administration is carried out using any type of formulation or dosage unit suitable for application to mucosal tissue. For example, the selected active agent may be administered to the buccal mucosa in an adhesive tablet or patch, sublingually administered by placing a solid dosage form under the tongue, lingually administered by placing a solid dosage form on the tongue, administered nasally as droplets or a nasal spray, administered by inhalation of an aerosol formulation, a non-aerosol liquid formulation, or a dry powder, placed within or near the rectum ("transrectal" formulations), or administered to the urethra as a suppository, ointment, or the like.

Preferred buccal dosage forms will typically comprise a therapeutically effective amount of an active agent and a bioerodible (hydrolyzable) polymeric carrier that may also serve to adhere the dosage form to the buccal mucosa. The buccal dosage unit is fabricated so as to erode over a predetermined time period, wherein drug delivery is provided essentially throughout. The time period is typically in the range of from about 1 hour to about 72 hours. Preferred buccal delivery preferably occurs over a time period of from about 2 hours to about 24 hours. Buccal drug delivery for short term use should preferably occur over a time period of from about 2 hours to about 8 hours, more preferably over a time period of from about 3 hours to about 4 hours. As needed buccal drug delivery preferably will occur over a time period of from about 1 hour to about 12 hours, more preferably from about 2 hours to about 8 hours, most preferably from about 3 hours to about 6 hours. Sustained buccal drug delivery will preferably occur over a time period of from about 6 hours to about 72 hours, more preferably from about 12 hours to about 48 hours, most preferably from about 24 hours to about 48 hours. Buccal drug delivery, as will be appreciated by those skilled in the art, avoids the disadvantages encountered with oral drug administration, e.g., slow absorption, degradation of the

active agent by fluids present in the gastrointestinal tract and/or first-pass inactivation in the liver.

The "therapeutically effective amount" of the active agent in the buccal dosage unit will of course depend on the potency of the agent and the intended dosage, which, in turn, is dependent on the particular individual undergoing treatment, the specific indication, and the like. The buccal dosage unit will generally contain from about 1.0 wt. % to about 60 wt. % active agent, preferably on the order of from about 1 wt. % to about 30 wt. % active agent. With regard to the bioerodible (hydrolyzable) polymeric carrier, it will be appreciated that virtually any such carrier can be used, so long as the desired drug release profile is not compromised, and the carrier is compatible with the active agents to be administered and any other components of the buccal dosage unit. Generally, the polymeric carrier comprises a hydrophilic (water-soluble and water-swellaable) polymer that adheres to the wet surface of the buccal mucosa. Examples of polymeric carriers useful herein include acrylic acid polymers and co, e.g., those known as "carbomers" (Carbopol®, which may be obtained from B. F. Goodrich, is one such polymer). Other suitable polymers include, but are not limited to: hydrolyzed polyvinylalcohol; polyethylene oxides (e.g., SENTRY Polyox® water soluble resins, available from Union Carbide); polyacrylates (e.g., Gantrez®, which may be obtained from GAF); vinyl polymers and copolymers; polyvinylpyrrolidone; dextran; guar gum; pectins; starches; and cellulosic polymers such as hydroxypropyl methylcellulose, (e.g., Methocel®, which may be obtained from the Dow Chemical Company), hydroxypropyl cellulose (e.g., Klucel®, which may also be obtained from Dow), hydroxypropyl cellulose ethers (see, e.g., U.S. Pat. No. 4,704,285 to Alderman), hydroxyethyl cellulose, carboxymethyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, ethyl cellulose, cellulose acetate phthalate, cellulose acetate butyrate, and the like.

Other components may also be incorporated into the buccal dosage forms described herein. The additional components include, but are not limited to, disintegrants, diluents, binders, lubricants, flavoring, colorants, preservatives, and the like. Examples of disintegrants that may be used include, but are not limited to, cross-linked polyvinylpyrrolidones, such as crospovidone (e.g., Polyplasdone® XL, which may be obtained from GAF), cross-linked carboxylic methylcelluloses, such as croscarmellose

(e.g., Ac-di-sol®, which may be obtained from FMC), alginic acid, and sodium carboxymethyl starches (e.g., Explotab®, which may be obtained from Edward Medall Co., Inc.), methylcellulose, agar bentonite and alginic acid. Suitable diluents are those which are generally useful in pharmaceutical formulations prepared using compression techniques, e.g., dicalcium phosphate dihydrate (e.g., Di-Tab®, which may be obtained from Stauffer), sugars that have been processed by cocrystallization with dextrin (e.g., co-crystallized sucrose and dextrin such as Di-Pak®, which may be obtained from Amstar), calcium phosphate, cellulose, kaolin, mannitol, sodium chloride, dry starch, powdered sugar and the like. Binders, if used, are those that enhance adhesion. Examples of such binders include, but are not limited to, starch, gelatin and sugars such as sucrose, dextrose, molasses, and lactose. Particularly preferred lubricants are stearates and stearic acid, and an optimal lubricant is magnesium stearate.

Sublingual and lingual dosage forms include tablets, creams, ointments, lozenges, pastes, and any other solid dosage form where the active ingredient is admixed into a disintegrable matrix. The tablet, cream, ointment or paste for sublingual or lingual delivery comprises a therapeutically effective amount of the selected active agent and one or more conventional nontoxic carriers suitable for sublingual or lingual drug administration. The sublingual and lingual dosage forms of the present invention can be manufactured using conventional processes. The sublingual and lingual dosage units are fabricated to disintegrate rapidly. The time period for complete disintegration of the dosage unit is typically in the range of from about 10 seconds to about 30 minutes, and optimally is less than 5 minutes.

Other components may also be incorporated into the sublingual and lingual dosage forms described herein. The additional components include, but are not limited to binders, disintegrants, wetting agents, lubricants, and the like. Examples of binders that may be used include water, ethanol, polyvinylpyrrolidone; starch solution gelatin solution, and the like. Suitable disintegrants include dry starch, calcium carbonate, polyoxyethylene sorbitan fatty acid esters, sodium lauryl sulfate, stearic monoglyceride, lactose, and the like. Wetting agents, if used, include glycerin, starches, and the like. Particularly preferred lubricants are stearates and polyethylene glycol. Additional components that may be incorporated into sublingual and lingual dosage forms are

known, or will be apparent, to those skilled in this art (See, e.g., Remington: The Science and Practice of Pharmacy, supra).

For transurethral administration, the formulation comprises a urethral dosage form containing the active agent and one or more selected carriers or excipients, such as water, silicone, waxes, petroleum jelly, polyethylene glycol ("PEG"), propylene glycol ("PG"), liposomes, sugars such as mannitol and lactose, and/or a variety of other materials, with polyethylene glycol and derivatives thereof particularly preferred.

Depending on the particular active agent administered, it may be desirable to incorporate a transurethral permeation enhancer in the urethral dosage form. Examples of suitable transurethral permeation enhancers include dimethylsulfoxide ("DMSO"), dimethyl formamide ("DMF"), N, N-dimethylacetamide ("DMA"), decylmethylsulfoxide ("C₁₀ MSO"), polyethylene glycol monolaurate ("PEGML"), glycerol monolaurate, lecithin, the 1-substituted azacycloheptan-2-ones, particularly 1-n-dodecylcyclazacycloheptan-2-one (available under the trademark Azone® from Nelson Research & Development Co., Irvine, Calif.), SEPA® (available from Macrochem Co., Lexington, Mass.), surfactants as discussed above, including, for example, Tergitol®, Nonoxynol-9® and TWEEN-80®, and lower alkanols such as ethanol.

Transurethral drug administration, as explained in U.S. Pat. Nos. 5,242,391, 5,474,535, 5,686,093 and 5,773,020, can be carried out in a number of different ways using a variety of urethral dosage forms. For example, the drug can be introduced into the urethra from a flexible tube, squeeze bottle, pump or aerosol spray. The drug may also be contained in coatings, pellets or suppositories that are absorbed, melted or bioeroded in the urethra. In certain embodiments, the drug is included in a coating on the exterior surface of a penile insert. It is preferred, although not essential, that the drug be delivered from at least about 3 cm into the urethra, and preferably from at least about 7 cm into the urethra. Generally, delivery from at least about 3 cm to about 8 cm into the urethra will provide effective results in conjunction with the present method.

Urethral suppository formulations containing PEG or a PEG derivative may be conveniently formulated using conventional techniques, e.g., compression molding, heat molding or the like, as will be appreciated by those skilled in the art and as described in the pertinent literature and pharmaceutical texts. (See, e.g., Remington: The Science and

Practice of Pharmacy, supra), which discloses typical methods of preparing pharmaceutical compositions in the form of urethral suppositories. The PEG or PEG derivative preferably has a molecular weight in the range of from about 200 to about 2,500 g/mol, more preferably in the range of from about 1,000 to about 2,000 g/mol. Suitable polyethylene glycol derivatives include polyethylene glycol fatty acid esters, for example, polyethylene glycol monostearate, polyethylene glycol sorbitan esters, e.g., polysorbates, and the like. Depending on the particular active agent, it may also be preferred that urethral suppositories contain one or more solubilizing agents effective to increase the solubility of the active agent in the PEG or other transurethral vehicle.

It may be desirable to deliver the active agent in a urethral dosage form that provides for controlled or sustained release of the agent. In such a case, the dosage form comprises a biocompatible, biodegradable material, typically a biodegradable polymer. Examples of such polymers include polyesters, polyalkylcyanoacrylates, polyorthoesters, polyanhydrides, albumin, gelatin and starch. As explained, for example, in PCT Publication No. WO 96/40054, these and other polymers can be used to provide biodegradable microparticles that enable controlled and sustained drug release, in turn minimizing the required dosing frequency.

The urethral dosage form will preferably comprise a suppository that is on the order of from about 2 to about 20 mm in length, preferably from about 5 to about 10 mm in length, and less than about 5 mm in width, preferably less than about 2 mm in width. The weight of the suppository will typically be in the range of from about 1 mg to about 100 mg, preferably in the range of from about 1 mg to about 50 mg. However, it will be appreciated by those skilled in the art that the size of the suppository can and will vary, depending on the potency of the drug, the nature of the formulation, and other factors.

Transurethral drug delivery may involve an "active" delivery mechanism such as iontophoresis, electroporation or phonophoresis. Devices and methods for delivering drugs in this way are well known in the art. Iontophoretically assisted drug delivery is, for example, described in PCT Publication No. WO 96/40054, cited above. Briefly, the active agent is driven through the urethral wall by means of an electric current passed from an external electrode to a second electrode contained within or affixed to a urethral probe.

Preferred transrectal dosage forms include rectal suppositories, creams, ointments, and liquid formulations (enemas). The suppository, cream, ointment or liquid formulation for transrectal delivery comprises a therapeutically effective amount of the selected phosphodiesterase inhibitor and one or more conventional nontoxic carriers suitable for transrectal drug administration. The transrectal dosage forms of the present invention can be manufactured using conventional processes. The transrectal dosage unit can be fabricated to disintegrate rapidly or over a period of several hours. The time period for complete disintegration is preferably in the range of from about 10 minutes to about 6 hours, and optimally is less than about 3 hours.

Other components may also be incorporated into the transrectal dosage forms described herein. The additional components include, but are not limited to, stiffening agents, antioxidants, preservatives, and the like. Examples of stiffening agents that may be used include, for example, paraffin, white wax and yellow wax. Preferred antioxidants, if used, include sodium bisulfite and sodium metabisulfite.

Preferred vaginal or perivaginal dosage forms include vaginal suppositories, creams, ointments, liquid formulations, pessaries, tampons, gels, pastes, foams or sprays. The suppository, cream, ointment, liquid formulation, pessary, tampon, gel, paste, foam or spray for vaginal or perivaginal delivery comprises a therapeutically effective amount of the selected active agent and one or more conventional nontoxic carriers suitable for vaginal or perivaginal drug administration. The vaginal or perivaginal forms of the present invention can be manufactured using conventional processes as disclosed in Remington: The Science and Practice of Pharmacy, supra (see also drug formulations as adapted in U.S. Patent Nos. 6,515,198; 6,500,822; 6,417,186; 6,416,779; 6,376,500; 6,355,641; 6,258,819; 6,172,062; and 6,086,909). The vaginal or perivaginal dosage unit can be fabricated to disintegrate rapidly or over a period of several hours. The time period for complete disintegration is preferably in the range of from about 10 minutes to about 6 hours, and optimally is less than about 3 hours.

Other components may also be incorporated into the vaginal or perivaginal dosage forms described herein. The additional components include, but are not limited to, stiffening agents, antioxidants, preservatives, and the like. Examples of stiffening

agents that may be used include, for example, paraffin, white wax and yellow wax. Preferred antioxidants, if used, include sodium bisulfite and sodium metabisulfite.

The active agents may also be administered intranasally or by inhalation. Compositions for intranasal administration are generally liquid formulations for administration as a spray or in the form of drops, although powder formulations for intranasal administration, e.g., insufflations, are also known, as are nasal gels, creams, pastes or ointments. For liquid formulations, the active agent can be formulated into a solution, e.g., water or isotonic saline, buffered or unbuffered, or as a suspension. Preferably, such solutions or suspensions are isotonic relative to nasal secretions and of about the same pH, ranging e.g., from about pH 4.0 to about pH 7.4 or, from about pH 6.0 to about pH 7.0. Buffers should be physiologically compatible and include, simply by way of example, phosphate buffers. Furthermore, various devices are available in the art for the generation of drops, droplets and sprays, including droppers, squeeze bottles, and manually and electrically powered intranasal pump dispensers. Active agent containing intranasal carriers may also include nasal gels, creams, pastes or ointments with a viscosity of, e.g., from about 10 to about 6500 cps, or greater, depending on the desired sustained contact with the nasal mucosal surfaces. Such carrier viscous formulations may be based upon, simply by way of example, alkylcelluloses and/or other biocompatible carriers of high viscosity well known to the art (see e.g., Remington: The Science and Practice of Pharmacy, supra). Other ingredients, such as art known preservatives, colorants, lubricating or viscous mineral or vegetable oils, perfumes, natural or synthetic plant extracts such as aromatic oils, and humectants and viscosity enhancers such as, e.g., glycerol, can also be included to provide additional viscosity, moisture retention and a pleasant texture and odor for the formulation. Formulations for inhalation may be prepared as an aerosol, either a solution aerosol in which the active agent is solubilized in a carrier (e.g., propellant) or a dispersion aerosol in which the active agent is suspended or dispersed throughout a carrier and an optional solvent. Non-aerosol formulations for inhalation may take the form of a liquid, typically an aqueous suspension, although aqueous solutions may be used as well. In such a case, the carrier is typically a sodium chloride solution having a concentration such that the formulation is isotonic relative to normal body fluid. In addition to the carrier, the liquid formulations

may contain water and/or excipients including an antimicrobial preservative (e.g., benzalkonium chloride, benzethonium chloride, chlorobutanol, phenylethyl alcohol, thimerosal and combinations thereof), a buffering agent (e.g., citric acid, potassium metaphosphate, potassium phosphate, sodium acetate, sodium citrate, and combinations thereof), a surfactant (e.g., polysorbate 80, sodium lauryl sulfate, sorbitan monopalmitate and combinations thereof), and/or a suspending agent (e.g., agar, bentonite, microcrystalline cellulose, sodium carboxymethylcellulose, hydroxypropyl methylcellulose, tragacanth, veegum and combinations thereof). Non-aerosol formulations for inhalation may also comprise dry powder formulations, particularly insufflations in which the powder has an average particle size of from about 0.1 μm to about 50 μm , preferably from about 1 μm to about 25 μm .

Topical Formulations

Topical formulations may be in any form suitable for application to the body surface, and may comprise, for example, an ointment, cream, gel, lotion, solution, paste or the like, and/or may be prepared so as to contain liposomes, micelles, and/or microspheres. Preferred topical formulations herein are ointments, creams and gels.

Ointments, as is well known in the art of pharmaceutical formulation, are semisolid preparations that are typically based on petrolatum or other petroleum derivatives. The specific ointment base to be used, as will be appreciated by those skilled in the art, is one that will provide for optimum drug delivery, and, preferably, will provide for other desired characteristics as well, e.g., emolliency or the like. As with other carriers or vehicles, an ointment base should be inert, stable, nonirritating and nonsensitizing. As explained in Remington: The Science and Practice of Pharmacy, supra, ointment bases may be grouped in four classes: oleaginous bases; emulsifiable bases; emulsion bases; and water-soluble bases. Oleaginous ointment bases include, for example, vegetable oils, fats obtained from animals, and semisolid hydrocarbons obtained from petroleum. Emulsifiable ointment bases, also known as absorbent ointment bases, contain little or no water and include, for example, hydroxystearin sulfate, anhydrous lanolin and hydrophilic petrolatum. Emulsion ointment bases are either water-in-oil (W/O) emulsions or oil-in-water (O/W) emulsions, and include, for

example, cetyl alcohol, glyceryl monostearate, lanolin and stearic acid. Preferred water-soluble ointment bases are prepared from polyethylene glycols of varying molecular weight (See, e.g., Remington: The Science and Practice of Pharmacy, supra).

Creams, as also well known in the art, are viscous liquids or semisolid emulsions, either oil-in-water or water-in-oil. Cream bases are water-washable, and contain an oil phase, an emulsifier and an aqueous phase. The oil phase, also called the "internal" phase, is generally comprised of petrolatum and a fatty alcohol such as cetyl or stearyl alcohol. The aqueous phase usually, although not necessarily, exceeds the oil phase in volume, and generally contains a humectant. The emulsifier in a cream formulation is generally a nonionic, anionic, cationic or amphoteric surfactant.

As will be appreciated by those working in the field of pharmaceutical formulation, gels are semisolid, suspension-type systems. Single-phase gels contain organic macromolecules distributed substantially uniformly throughout the carrier liquid, which is typically aqueous, but also, preferably, contain an alcohol and, optionally, an oil. Preferred "organic macromolecules," i.e., gelling agents, are crosslinked acrylic acid polymers such as the "carbomer" family of polymers, e.g., carboxypolyalkylenes that may be obtained commercially under the Carbopol® trademark. Also preferred are hydrophilic polymers such as polyethylene oxides, polyoxyethylene-polyoxypropylene copolymers and polyvinylalcohol; cellulosic polymers such as hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose phthalate, and methylcellulose; gums such as tragacanth and xanthan gum; sodium alginate; and gelatin. In order to prepare a uniform gel, dispersing agents such as alcohol or glycerin can be added, or the gelling agent can be dispersed by trituration, mechanical mixing, and/or stirring.

Various additives, known to those skilled in the art, may be included in the topical formulations. For example, solubilizers may be used to solubilize certain active agents. For those drugs having an unusually low rate of permeation through the skin or mucosal tissue, it may be desirable to include a permeation enhancer in the formulation; suitable enhancers are as described elsewhere herein.

Transdermal Administration

The compounds of the invention may also be administered through the skin or mucosal tissue using conventional transdermal drug delivery systems, wherein the agent is contained within a laminated structure (typically referred to as a transdermal "patch") that serves as a drug delivery device to be affixed to the skin. Transdermal drug delivery may involve passive diffusion or it may be facilitated using electrotransport, e.g., iontophoresis. In a typical transdermal "patch," the drug composition is contained in a layer, or "reservoir," underlying an upper backing layer. The laminated structure may contain a single reservoir, or it may contain multiple reservoirs. In one type of patch, referred to as a "monolithic" system, the reservoir is comprised of a polymeric matrix of a pharmaceutically acceptable contact adhesive material that serves to affix the system to the skin during drug delivery. Examples of suitable skin contact adhesive materials include, but are not limited to, polyethylenes, polysiloxanes, polyisobutylenes, polyacrylates, polyurethanes, and the like. Alternatively, the drug-containing reservoir and skin contact adhesive are separate and distinct layers, with the adhesive underlying the reservoir which, in this case, may be either a polymeric matrix as described above, or it may be a liquid or hydrogel reservoir, or may take some other form.

The backing layer in these laminates, which serves as the upper surface of the device, functions as the primary structural element of the laminated structure and provides the device with much of its flexibility. The material selected for the backing material should be selected so that it is substantially impermeable to the active agent and any other materials that are present, the backing is preferably made of a sheet or film of a flexible elastomeric material. Examples of polymers that are suitable for the backing layer include polyethylene, polypropylene, polyesters, and the like.

During storage and prior to use, the laminated structure includes a release liner. Immediately prior to use, this layer is removed from the device to expose the basal surface thereof, either the drug reservoir or a separate contact adhesive layer, so that the system may be affixed to the skin. The release liner should be made from a drug/vehicle impermeable material.

Transdermal drug delivery systems may in addition contain a skin permeation enhancer. That is, because the inherent permeability of the skin to some drugs may be too low to allow therapeutic levels of the drug to pass through a reasonably sized area of

unbroken skin, it is necessary to coadminister a skin permeation enhancer with such drugs. Suitable enhancers are well known in the art and include, for example, those enhancers listed above in transmucosal compositions.

Parenteral Administration

Parenteral administration, if used, is generally characterized by injection, including intramuscular, intraperitoneal, intravenous (IV) and subcutaneous injection. Injectable formulations can be prepared in conventional forms, either as liquid solutions or suspensions; solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Preferably, sterile injectable suspensions are formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable formulation may also be a sterile injectable solution or a suspension in a nontoxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. A more recently revised approach for parenteral administration involves use of a slow release or sustained release system (See, e.g., U.S. Pat. No. 3,710,795).

Intravesical Administration

Intravesical administration, if used, is generally characterized by administration directly into the bladder and may include methods as described elsewhere herein. Other methods of intravesical administration may include those described in U.S. Patent Nos. 6,207,180 and 6,039,967, as well as other methods that are known to one of skill in the art.

Intrathecal Administration

Intrathecal administration, if used, is generally characterized by administration directly into the intrathecal space (where fluid flows around the spinal cord).

One common system utilized for intrathecal administration is the APT Intrathecal treatment system available from Medtronic, Inc. APT Intrathecal uses a small pump that

is surgically placed under the skin of the abdomen to deliver medication directly into the intrathecal space. The medication is delivered through a small tube called a catheter that is also surgically placed.

Another system available from Medtronic that is commonly utilized for intrathecal administration is the fully implantable, programmable SynchroMed[®] Infusion System. The SynchroMed[®] Infusion System has two parts that are both placed in the body during a surgical procedure: the catheter and the pump. The catheter is a small, soft tube. One end is connected to the catheter port of the pump, and the other end is placed in the intrathecal space. The pump is a round metal device about one inch (2.5 cm) thick, three inches (8.5 cm) in diameter, and weighs about six ounces (205 g) that stores and releases prescribed amounts of medication directly into the intrathecal space. It is made of titanium, a lightweight, medical-grade metal. The reservoir is the space inside the pump that holds the medication. The fill port is a raised center portion of the pump through which the pump is refilled. The doctor or a nurse inserts a needle through the patient's skin and through the fill port to fill the pump. Some pumps have a side catheter access port that allows the doctor to inject other medications or sterile solutions directly into the catheter, bypassing the pump.

The SynchroMed[®] pump automatically delivers a controlled amount of medication through the catheter to the intrathecal space around the spinal cord, where it is most effective. The exact dosage, rate and timing prescribed by the doctor are entered in the pump using a programmer, an external computer-like device that controls the pump's memory. Information about the patient's prescription is stored in the pump's memory. The doctor can easily review this information by using the programmer. The programmer communicates with the pump by radio signals that allow the doctor to tell how the pump is operating at any given time. The doctor also can use the programmer to change your medication dosage.

Methods of intrathecal administration may include those described above available from Medtronic, as well as other methods that are known to one of skill in the art.

Additional Dosage Formulations and Drug Delivery Systems

As compared with traditional drug delivery approaches, some controlled release technologies rely upon the modification of both macromolecules and synthetic small molecules to allow them to be actively instead of passively absorbed into the body. For example, XenoPort Inc. utilizes technology that takes existing molecules and re-engineers them to create new chemical entities (unique molecules) that have improved pharmacologic properties to either: 1) lengthen the short half-life of a drug; 2) overcome poor absorption; and/or 3) deal with poor drug distribution to target tissues. Techniques to lengthen the short half-life of a drug include the use of prodrugs with slow cleavage rates to release drugs over time or that engage transporters in small and large intestines to allow the use of oral sustained delivery systems, as well as drugs that engage active transport systems. Examples of such controlled release formulations, tablets, dosage forms, and drug delivery systems, and that are suitable for use with the present invention, are described in the following published US and PCT patent applications assigned to Xenoport Inc.: US20030158254; US20030158089; US20030017964; US2003130246; WO02100172; WO02100392; WO02100347; WO02100344; WO0242414; WO0228881; WO0228882; WO0244324; WO0232376; WO0228883; and WO0228411. In particular, Xenoport's XP13512 is a transported Prodrug of gabapentin that has been engineered to utilize high capacity transport mechanisms located in both the small and large intestine and to rapidly convert to gabapentin once in the body. In contrast to gabapentin itself, XP13512 was shown in preclinical and clinical studies to produce dose proportional blood levels of gabapentin across a broad range of oral doses, and to be absorbed efficiently from the large intestine.

Some other controlled release technologies rely upon methods that promote or enhance gastric retention, such as those developed by Depomed Inc. Because many drugs are best absorbed in the stomach and upper portions of the small intestine, Depomed has developed tablets that swell in the stomach during the postprandial or fed mode so that they are treated like undigested food. These tablets therefore sit safely and neutrally in the stomach for 6, 8, or more hours and deliver drug at a desired rate and time to upper gastrointestinal sites. Specific technologies in this area include: 1) tablets that slowly erode in gastric fluids to deliver drugs at almost a constant rate (particularly useful

for highly insoluble drugs); 2) bi-layer tablets that combine drugs with different characteristics into a single table (such as a highly insoluble drug in an erosion layer and a soluble drug in a diffusion layer for sustained release of both); and 3) combination tablets that can either deliver drugs simultaneously or in sequence over a desired period of time (including an initial burst of a fast acting drug followed by slow and sustained delivery of another drug). Examples of such controlled release formulations that are suitable for use with the present invention and that rely upon gastric retention during the postprandial or fed mode, include tablets, dosage forms, and drug delivery systems in the following US patents assigned to Depomed Inc.: US 6,488,962; US 6,451,808; US 6,340,475; US 5,972,389; US 5,582,837; and US 5,007,790. Examples of such controlled release formulations that are suitable for use with the present invention and that rely upon gastric retention during the postprandial or fed mode, include tablets, dosage forms, and drug delivery systems in the following published US and PCT patent applications assigned to Depomed Inc.: US20030147952; US20030104062; US20030104053; US20030104052; US20030091630; US20030044466; US20030039688; US20020051820; WO0335040; WO0335039; WO0156544; WO0132217; WO9855107; WO9747285; and WO9318755.

Other controlled release systems include those developed by ALZA Corporation based upon: 1) osmotic technology for oral delivery; 2) transdermal delivery via patches; 3) liposomal delivery via intravenous injection; 4) osmotic technology for long-term delivery via implants; and 5) depot technology designed to deliver agents for periods of days to a month. ALZA oral delivery systems include those that employ osmosis to provide precise, controlled drug delivery for up to 24 hours for both poorly soluble and highly soluble drugs, as well as those that deliver high drug doses meeting high drug loading requirements. ALZA controlled transdermal delivery systems provide drug delivery through intact skin for as long as one week with a single application to improve drug absorption and deliver constant amounts of drug into the bloodstream over time. ALZA liposomal delivery systems involve lipid nanoparticles that evade recognition by the immune system because of their unique polyethylene glycol (PEG) coating, allowing the precise delivery of drugs to disease-specific areas of the body. ALZA also has developed osmotically driven systems to enable the continuous delivery of small drugs,

peptides, proteins, DNA and other bioactive macromolecules for up to one year for systemic or tissue-specific therapy. Finally, ALZA depot injection therapy is designed to deliver biopharmaceutical agents and small molecules for periods of days to a month using a nonaqueous polymer solution for the stabilization of macromolecules and a unique delivery profile.

Examples of controlled release formulations, tablets, dosage forms, and drug delivery systems that are suitable for use with the present invention are described in the following US patents assigned to ALZA Corporation: US 4,367,741; US 4,402,695; US 4,418,038; US 4,434,153; US 4,439,199; US 4,450,198; US 4,455,142; US 4,455,144; US 4,484,923; US 4,486,193; US 4,489,197; US 4,511,353; US 4,519,801; US 4,526,578; US 4,526,933; US 4,534,757; US 4,553,973; US 4,559,222; US 4,564,364; US 4,578,075; US 4,588,580; US 4,610,686; US 4,618,487; US 4,627,851; US 4,629,449; US 4,642,233; US 4,649,043; US 4,650,484; US 4,659,558; US 4,661,105; US 4,662,880; US 4,675,174; US 4,681,583; US 4,684,524; US 4,692,336; US 4,693,895; US 4,704,119; US 4,705,515; US 4,717,566; US 4,721,613; US 4,723,957; US 4,725,272; US 4,728,498; US 4,743,248; US 4,747,847; US 4,751,071; US 4,753,802; US 4,755,180; US 4,756,314; US 4,764,380; US 4,773,907; US 4,777,049; US 4,781,924; US 4,786,503; US 4,788,062; US 4,810,502; US 4,812,313; US 4,816,258; US 4,824,675; US 4,834,979; US 4,837,027; US 4,842,867; US 4,846,826; US 4,847,093; US 4,849,226; US 4,851,229; US 4,851,231; US 4,851,232; US 4,853,229; US 4,857,330; US 4,859,470; US 4,863,456; US 4,863,744; US 4,865,598; US 4,867,969; US 4,871,548; US 4,872,873; US 4,874,388; US 4,876,093; US 4,892,778; US 4,902,514; US 4,904,474; US 4,913,903; US 4,915,949; US 4,915,952; US 4,917,895; US 4,931,285; US 4,946,685; US 4,948,592; US 4,954,344; US 4,957,494; US 4,960,416; US 4,961,931; US 4,961,932; US 4,963,141; US 4,966,769; US 4,971,790; US 4,976,966; US 4,986,987; US 5,006,346; US 5,017,381; US 5,019,397; US 5,023,076; US 5,023,088; US 5,024,842; US 5,028,434; US 5,030,454; US 5,071,656; US 5,077,054; US 5,082,668; US 5,104,390; US 5,110,597; US 5,122,128; US 5,125,894; US 5,141,750; US 5,141,752; US 5,156,850; US 5,160,743; US 5,160,744; US 5,169,382; US 5,171,576; US 5,176,665; US 5,185,158; US 5,190,765; US 5,198,223; US 5,198,229; US 5,200,195; US 5,200,196; US 5,204,116;

US 5,208,037; US 5,209,746; US 5,221,254; US 5,221,278; US 5,229,133; US
5,232,438; US 5,232,705; US 5,236,689; US 5,236,714; US 5,240,713; US 5,246,710;
US 5,246,711; US 5,252,338; US 5,254,349; US 5,266,332; US 5,273,752; US
5,284,660; US 5,286,491; US 5,308,348; US 5,318,558; US 5,320,850; US 5,322,502;
US 5,326,571; US 5,330,762; US 5,338,550; US 5,340,590; US 5,342,623; US
5,344,656; US 5,348,746; US 5,358,721; US 5,364,630; US 5,376,377; US 5,391,381;
US 5,402,777; US 5,403,275; US 5,411,740; US 5,417,675; US 5,417,676; US
5,417,682; US 5,423,739; US 5,424,289; US 5,431,919; US 5,443,442; US 5,443,459;
US 5,443,461; US 5,456,679; US 5,460,826; US 5,462,741; US 5,462,745; US
5,489,281; US 5,499,979; US 5,500,222; US 5,512,293; US 5,512,299; US 5,529,787;
US 5,531,736; US 5,532,003; US 5,533,971; US 5,534,263; US 5,540,912; US
5,543,156; US 5,571,525; US 5,573,503; US 5,591,124; US 5,593,695; US 5,595,759;
US 5,603,954; US 5,607,696; US 5,609,885; US 5,614,211; US 5,614,578; US
5,620,705; US 5,620,708; US 5,622,530; US 5,622,944; US 5,633,011; US 5,639,477;
US 5,660,861; US 5,667,804; US 5,667,805; US 5,674,895; US 5,688,518; US
5,698,224; US 5,702,725; US 5,702,727; US 5,707,663; US 5,713,852; US 5,718,700;
US 5,736,580; US 5,770,227; US 5,780,058; US 5,783,213; US 5,785,994; US
5,795,591; US 5,811,465; US 5,817,624; US 5,824,340; US 5,830,501; US 5,830,502;
US 5,840,754; US 5,858,407; US 5,861,439; US 5,863,558; US 5,876,750; US
5,883,135; US 5,897,878; US 5,904,934; US 5,904,935; US 5,906,832; US 5,912,268;
US 5,914,131; US 5,916,582; US 5,932,547; US 5,938,654; US 5,941,844; US
5,955,103; US 5,972,369; US 5,972,370; US 5,972,379; US 5,980,943; US 5,981,489;
US 5,983,130; US 5,989,590; US 5,995,869; US 5,997,902; US 6,001,390; US
6,004,309; US 6,004,578; US 6,008,187; US 6,020,000; US 6,034,101; US 6,036,973;
US 6,039,977; US 6,057,374; US 6,066,619; US 6,068,850; US 6,077,538; US
6,083,190; US 6,096,339; US 6,106,845; US 6,110,499; US 6,120,798; US 6,120,803;
US 6,124,261; US 6,130,200; US 6,146,662; US 6,153,678; US 6,174,547; US
6,183,466; US 6,203,817; US 6,210,712; US 6,210,713; US 6,224,907; US 6,235,712;
US 6,245,357; US 6,262,115; US 6,264,990; US 6,267,984; US 6,287,598; US
6,289,241; US 6,331,311; US 6,333,050; US 6,342,249; US 6,346,270; US 6,365,183; US
6,368,626; US 6,387,403; US 6,419,952; US 6,440,457; US 6,468,961; US 6,491,683;

US 6,512,010; US 6,514,530; US 6,534,089; US 6,544,252; US 6,548,083; US 6,551,613; US 6,572,879; and US 6,596,314.

Other examples of controlled release formulations, tablets, dosage forms, and drug delivery systems that are suitable for use with the present invention are described in the following published US patent application and PCT applications assigned to ALZA Corporation: US20010051183; WO0004886; WO0013663; WO0013674; WO0025753; WO0025790; WO0035419; WO0038650; WO0040218; WO0045790; WO0066126; WO0074650; WO0119337; WO0119352; WO0121211; WO0137815; WO0141742; WO0143721; WO0156543; WO3041684; WO3041685; WO3041757; WO3045352; WO3051341; WO3053400; WO3053401; WO9000416; WO9004965; WO9113613; WO9116884; WO9204011; WO9211843; WO9212692; WO9213521; WO9217239; WO9218102; WO9300071; WO9305843; WO9306819; WO9314813; WO9319739; WO9320127; WO9320134; WO9407562; WO9408572; WO9416699; WO9421262; WO9427587; WO9427589; WO9503823; WO9519174; WO9529665; WO9600065; WO9613248; WO9625922; WO9637202; WO9640049; WO9640050; WO9640139; WO9640364; WO9640365; WO9703634; WO9800158; WO9802169; WO9814168; WO9816250; WO9817315; WO9827962; WO9827963; WO9843611; WO9907342; WO9912526; WO9912527; WO9918159; WO9929297; WO9929348; WO9932096; WO9932153; WO9948494; WO9956730; WO9958115; and WO9962496.

Another drug delivery technology suitable for use in the present invention is that disclosed by DepoMed, Inc. in U.S. Patent No. 6,682,759, which discloses a method for manufacturing a pharmaceutical tablet for oral administration combining both immediate-release and prolonged-release modes of drug delivery. The tablet according to the method comprises a prolonged-release drug core and an immediate-release drug coating or layer, which can be insoluble or sparingly soluble in water. The method limits the drug particle diameter in the immediate-release coating or layer to 10 microns or less. The coating or layer is either the particles themselves, applied as an aqueous suspension, or a solid composition that contains the drug particles incorporated in a solid material that disintegrates rapidly in gastric fluid.

Andrx Corporation has also developed drug delivery technology suitable for use in the present invention that includes: 1) a pelletized pulsatile delivery system ("PPDS");

2) a single composition osmotic tablet system ("SCOT"); 3) a solubility modulating hydrogel system ("SMHS"); 4) a delayed pulsatile hydrogel system ("DPHS"); 5) a stabilized pellet delivery system ("SPDS"); 6) a granulated modulating hydrogel system ("GMHS"); 7) a pelletized tablet system ("PELTAB"); 8) a porous tablet system ("PORTAB"); and 9) a stabilized tablet delivery system ("STDS"). PPDS uses pellets that are coated with specific polymers and agents to control the release rate of the microencapsulated drug and is designed for use with drugs that require a pulsed release. SCOT utilizes various osmotic modulating agents as well as polymer coatings to provide a zero-order drug release. SMHS utilizes a hydrogel-based dosage system that avoids the "initial burst effect" commonly observed with other sustained-release hydrogel formulations and that provides for sustained release without the need to use special coatings or structures that add to the cost of manufacturing. DPHS is designed for use with hydrogel matrix products characterized by an initial zero-order drug release followed by a rapid release that is achieved by the blending of selected hydrogel polymers to achieve a delayed pulse. SPDS incorporates a pellet core of drug and protective polymer outer layer, and is designed specifically for unstable drugs, while GMHS incorporates hydrogel and binding polymers with the drug and forms granules that are pressed into tablet form. PELTAB provides controlled release by using a water insoluble polymer to coat discrete drug crystals or pellets to enable them to resist the action of fluids in the gastrointestinal tract, and these coated pellets are then compressed into tablets. PORTAB provides controlled release by incorporating an osmotic core with a continuous polymer coating and a water soluble component that expands the core and creates microporous channels through which drug is released. Finally, STDS includes a dual layer coating technique that avoids the need to use a coating layer to separate the enteric coating layer from the omeprazole core.

Examples of controlled release formulations, tablets, dosage forms, and drug delivery systems that are suitable for use with the present invention are described in the following US patents assigned to Andrx Corporation: US 5,397,574; US 5,419,917; US 5,458,887; US 5,458,888; US 5,472,708; US 5,508,040; US 5,558,879; US 5,567,441; US 5,654,005; US 5,728,402; US 5,736,159; US 5,830,503; US 5,834,023; US 5,837,379; US 5,916,595; US 5,922,352; US 6,099,859; US 6,099,862; US 6,103,263;

US 6,106,862; US 6,156,342; US 6,177,102; US 6,197,347; US 6,210,716; US 6,238,703; US 6,270,805; US 6,284,275; US 6,485,748; US 6,495,162; US 6,524,620; US 6,544,556; US 6,589,553; US 6,602,522; and US 6,610,326.

Examples of controlled release formulations, tablets, dosage forms, and drug delivery systems that are suitable for use with the present invention are described in the following published US and PCT patent applications assigned to Andrx Corporation: US20010024659; US20020115718; US20020156066; WO0004883; WO0009091; WO0012097; WO0027370; WO0050010; WO0132161; WO0134123; WO0236077; WO0236100; WO02062299; WO02062824; WO02065991; WO02069888; WO02074285; WO03000177; WO9521607; WO9629992; WO9633700; WO9640080; WO9748386; WO9833488; WO9833489; WO9930692; WO9947125; and WO9961005.

Some other examples of drug delivery approaches focus on non-oral drug delivery, providing parenteral, transmucosal, and topical delivery of proteins, peptides, and small molecules. For example, the Atrigel[®] drug delivery system marketed by Atrix Laboratories Inc. comprises biodegradable polymers, similar to those used in biodegradable sutures, dissolved in biocompatible carriers. These pharmaceuticals may be blended into a liquid delivery system at the time of manufacturing or, depending upon the product, may be added later by a physician at the time of use. Injection of the liquid product subcutaneously or intramuscularly through a small gauge needle, or placement into accessible tissue sites through a cannula, causes displacement of the carrier with water in the tissue fluids, and a subsequent precipitate to form from the polymer into a solid film or implant. The drug encapsulated within the implant is then released in a controlled manner as the polymer matrix biodegrades over a period ranging from days to months. Examples of such drug delivery systems include Atrix's Eligard[®], Atridox[®]/Doxirobe[®], Atrisorb[®] FreeFlow[™]/Atrisorb[®]-D FreeFlow, bone growth products, and others as described in the following published US and PCT patent applications assigned to Atrix Laboratories Inc.: US RE37950; US 6,630,155; US 6,566,144; US 6,610,252; US 6,565,874; US 6,528,080; US 6,461,631; US 6,395,293; US 6,261,583; US 6,143,314; US 6,120,789; US 6,071,530; US 5,990,194; US 5,945,115; US 5,888,533; US 5,792,469; US 5,780,044; US 5,759,563; US 5,744,153; US 5,739,176; US 5,736,152; US 5,733,950; US 5,702,716; US 5,681,873; US 5,660,849; US 5,599,552;

US 5,487,897; US 5,368,859; US 5,340,849; US 5,324,519; US 5,278,202; US 5,278,201; US20020114737, US20030195489; US20030133964; US 20010042317; US20020090398; US20020001608; and US2001042317.

Atrix Laboratories Inc. also markets technology for the non-oral transmucosal delivery of drugs over a time period from minutes to hours. For example, Atrix's BEMA™ (Bioerodible Muco-Adhesive Disc) drug delivery system comprises pre-formed bioerodible discs for local or systemic delivery. Examples of such drug delivery systems include those as described in US Patent No. 6,245,345.

Other drug delivery systems marketed by Atrix Laboratories Inc. focus on topical drug delivery. For example, SMP™ (Solvent Particle System) allows the topical delivery of highly water-insoluble drugs. This product allows for a controlled amount of a dissolved drug to permeate the epidermal layer of the skin by combining the dissolved drug with a microparticle suspension of the drug. The SMP™ system works in stages whereby: 1) the product is applied to the skin surface; 2) the product near follicles concentrates at the skin pore; 3) the drug readily partitions into skin oils; and 4) the drug diffuses throughout the area. By contrast, MCA® (Mucocutaneous Absorption System) is a water-resistant topical gel providing sustained drug delivery. MCA® forms a tenacious film for either wet or dry surfaces where: 1) the product is applied to the skin or mucosal surface; 2) the product forms a tenacious moisture-resistant film; and 3) the adhered film provides sustained release of drug for a period from hours to days. Yet another product, BCP™ (Biocompatible Polymer System) provides a non-cytotoxic gel or liquid that is applied as a protective film for wound healing. Examples of these systems include Orajel®-Ultra Mouth Sore Medicine as well as those as described in the following published US patents and applications assigned to Atrix Laboratories Inc.: US 6,537,565; US 6,432,415; US 6,355,657; US 5,962,006; US 5,725,491; US 5,722,950; US 5,717,030; US 5,707,647; US 5,632,727; and US20010033853.

Additional formulations and compositions available from Teva Pharmaceutical Industries Ltd., Warner Lambert & Co., and Godecke Aktiengesellschaft that include gabapentin and are useful in the present invention include those as described in the following US patents and published US and PCT patent applications: US 6,531,509; US

6,255,526; US 6,054,482; US2003055109; US2002045662; US2002009115; WO 01/97782; WO 01/97612; EP 2001946364; WO 99/59573; WO 99/59572.

Dosage and Administration

The concentration of the active agent in any of the aforementioned dosage forms and compositions can vary a great deal, and will depend on a variety of factors, including the type of composition or dosage form, the corresponding mode of administration, the nature and activity of the specific active agent, and the intended drug release profile. Preferred dosage forms contain a unit dose of active agent, i.e., a single therapeutically effective dose. For creams, ointments, etc., a "unit dose" requires an active agent concentration that provides a unit dose in a specified quantity of the formulation to be applied. The unit dose of any particular active agent will depend, of course, on the active agent and on the mode of administration.

For the active agents of the present invention (including an $\alpha_2\delta$ subunit calcium channel modulator in combination with a compound with smooth muscle modulatory effects), the unit dose for oral, transmucosal, topical, transdermal, and parenteral administration will be in the range of from about 1 ng to about 10,000 mg, typically in the range of from about 100 ng to about 5,000 mg. Alternatively, for active agents of the present invention (including an $\alpha_2\delta$ subunit calcium channel modulator in combination with a compound with smooth muscle modulatory effects), the unit dose for oral, transmucosal, topical, transdermal, and parenteral administration will be greater than about 1 ng, about 5 ng, about 10 ng, about 20 ng, about 30 ng, about 40 ng, about 50 ng, about 100 ng, about 200 ng, about 300 ng, about 400 ng, about 500 ng, about 1 μ g, about 5 μ g, about 10 μ g, about 20 μ g, about 30 μ g, about 40 μ g, about 50 μ g, about 100 μ g, about 200 μ g, about 300 μ g, about 400 μ g, about 500 μ g, about .5 mg, about 1 mg, about 1.5 mg, about 2.0 mg, about 2.5 mg, about 3.0 mg, about 3.5 mg, about 4.0 mg, about 4.5 mg, about 5 mg, about 10 mg, about 20 mg, about 30 mg, about 40 mg, about 50 mg, about 100 mg, about 200 mg, about 300 mg, about 400 mg, about 500 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1025 mg, about 1050

mg, about 1075 mg, about 1100 mg, about 1125 mg, about 1150 mg, about 1175 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg, about 1975 mg, about 2000 mg, about 2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225 mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about 2400 mg, about 2425 mg, about 2450 mg, about 2475 mg, about 2500 mg, about 2525 mg, about 2550 mg, about 2575 mg, about 2600 mg, about 3,000 mg, about 3,500 mg, about 4,000 mg, about 4,500 mg, about 5,000 mg, about 5,500 mg, about 6,000 mg, about 6,500 mg, about 7,000 mg, about 7,500 mg, about 8,000 mg, about 8,500 mg, about 9,000 mg, or about 9,500 mg.

For the active agents of the present invention (including an $\alpha_2\delta$ subunit calcium channel modulator in combination with a compound with smooth muscle modulatory effects), the unit dose for intrathecal administration will be in the range of from about 1 fg to about 1 mg, typically in the range of from about 100 fg to about 1 ng. Alternatively, for the active agents of the present invention (including an $\alpha_2\delta$ subunit calcium channel modulator in combination with a compound with smooth muscle modulatory effects), the unit dose for intrathecal administration will be greater than about 1 fg, about 5 fg, about 10 fg, about 20 fg, about 30 fg, about 40 fg, about 50 fg, about 100 fg, about 200 fg, about 300 fg, about 400 fg, about 500 fg, about 1 pg, about 5 pg, about 10 pg, about 20 pg, about 30 pg, about 40 pg, about 50 pg, about 100 pg, about 200 pg, about 300 pg, about 400 pg, about 500 pg, about 1 ng, about 5 ng, about 10 ng, about 20 ng, about 30 ng, about 40 ng, about 50 ng, about 100 ng, about 200 ng, about 300 ng, about 400 ng, about 500 ng, about 1 μ g, about 5 μ g, about 10 μ g, about 20 μ g, about 30 μ g, about 40 μ g, about 50 μ g, about 100 μ g, about 200 μ g, about 300 μ g, about 400 μ g, or about 500 μ g.

The present invention also encompasses a pharmaceutical formulation encompassing an oral immediate release dosage form of oxybutynin in combination with gabapentin, wherein the unit dose of said oxybutynin will be in an amount less than about 5 mg, less than about 4.5 mg, less than about 4 mg, less than about 3.5 mg, less than about 3 mg, less than about 2.5 mg, less than about 2 mg, less than about 1.5 mg, or less than about .5 mg, and wherein the unit dose of said gabapentin will be in an amount between about 100 mg and about 300 mg, between about 125 mg and about 275 mg, between about 150 mg and about 250 mg, between about 175 mg and about 225 mg, or about 200 mg.

A therapeutically effective amount of a particular active agent administered to a given individual will, of course, be dependent on a number of factors, including the concentration of the specific active agent, composition or dosage form, the selected mode of administration, the age and general condition of the individual being treated, the sex of the individual, the severity of the individual's condition, and other factors known to the prescribing physician.

A therapeutically effective amount according to the present invention can include an amount effective to show a reduction in the frequency or intensity of at least one symptom associated with the functional bowel disorder. For example, in a patient with IBS, an effective amount of the active agents of the present invention would be an amount effective at reducing abdominal pain or discomfort associated with bowel movements. A further example of an effective amount of the active agents of the present invention for treating a patient with IBS would be an amount effective for reducing the onset or frequency of loose stool.

Packaged Kits

In another embodiment, a packaged kit is provided that contains the pharmaceutical formulation to be administered, i.e., a pharmaceutical formulation containing a therapeutically effective amount of an $\alpha_2\delta$ subunit calcium channel modulator in combination with one or more compounds with smooth muscle modulatory effects for the treatment of functional bowel disorders, particularly IBS, a container, preferably sealed, for housing the formulation during storage and prior to use, and

instructions for carrying out drug administration in a manner effective to treat functional bowel disorders, particularly IBS. The instructions will typically be written instructions on a package insert and/or on a label. Depending on the type of formulation and the intended mode of administration, the kit may also include a device for administering the formulation. Formulations may be any suitable formulations as described herein. For example, formulations may be an oral dosage form containing a unit dosage of a selected active agent.

The kit may contain multiple formulations of different dosages of the same agent. The kit may also contain multiple formulations of different active agents. The kit may contain formulations suitable for sequential, separate and/or simultaneous use in the treatment of lower urinary tract disorders, and instructions for carrying out drug administration where the formulations are administered sequentially, separately and/or simultaneously in the treatment of lower urinary tract disorders.

The parts of the kit may be independently held in one or more containers--such as bottles, syringes, plates, wells, blister packs etc.

Insurance Claims

In general, the processing of an insurance claim for the coverage of a given medical treatment or drug therapy involves notification of the insurance company, or any other entity, that has issued the insurance policy against which the claim is being filed, that the medical treatment or drug therapy will be performed. A determination is then made as to whether the medical treatment or drug therapy that will be performed is covered under the terms of the policy. If covered, the claim is then processed, which can include payment, reimbursement, or application against a deductible.

The present invention encompasses a method for processing an insurance claim under an insurance policy for an $\alpha_2\delta$ subunit calcium channel modulator and an antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof used in the treatment functional bowel disorders, particularly IBS, wherein said $\alpha_2\delta$ subunit calcium channel modulator and antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof are administered sequentially or concurrently in different compositions. This method

comprises: 1) receiving notification that treatment using said $\alpha_2\delta$ subunit calcium channel modulator and said antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs or active metabolites thereof will be performed or notification of a prescription; 2) determining whether said treatment using said $\alpha_2\delta$ subunit calcium channel modulator and said antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs or active metabolites is covered under said insurance policy; and 3) processing said claim for treatment of said functional bowel disorders using said $\alpha_2\delta$ subunit calcium channel modulator and said antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof, including payment, reimbursement, or application against a deductible. For use in this method, a particularly preferred $\alpha_2\delta$ subunit calcium channel modulator is gabapentin, while a particularly preferred antimuscarinic is oxybutynin. This method also encompasses the processing of claims for and $\alpha_2\delta$ subunit calcium channel modulator, particularly gabapentin, or an antimuscarinic, particularly oxybutynin, when either has been prescribed separately or concurrently for the treatment of functional bowel disorders, particularly IBS.

Many modifications and other embodiments of the inventions set forth herein will come to mind to one skilled in the art to which these inventions pertain having the benefit of the teachings presented in the foregoing descriptions and the associated drawings. Therefore, it is to be understood that the inventions are not to be limited to the specific embodiments disclosed and that modifications and other embodiments are intended to be included within the scope of the appended embodiments. Although specific terms are employed herein, they are used in a generic and descriptive sense only and not for purposes of limitation.

EXAMPLES

Methods For Treating Functional Bowel Disorders Using $\alpha_2\delta$ Subunit Calcium Channel Modulators With Smooth Muscle Modulators

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims. The following examples

illustrate the effects of administration of the combination of an $\alpha_2\delta$ subunit calcium channel modulator and a smooth muscle modulator in various well-accepted models of functional bowel disorders. It is expected that these results will demonstrate the efficacy of the combination of an $\alpha_2\delta$ subunit calcium channel modulator and a smooth muscle modulator for treatment of functional bowel disorders as described herein.

DISTENSION MODELS

A variety of assays can be used to assess visceromotor and pain responses to rectal distension. See, for example, Gunter *et al.*, *Physiol. Behav.*, 69(3): 379-82 (2000), Depoortere *et al.*, *J. Pharmacol. and Exp. Ther.*, 294(3): 983-990 (2000), Morteau *et al.*, *Fund. Clin. Pharmacol.*, 8(6): 553-62 (1994), Gibson *et al.*, *Gastroenterology (Suppl. 1)*, 120(5): A19-A20 (2001) and Gschossmann *et al.*, *Eur. J. Gastro. Hepat.*, 14(10): 1067-72 (2002) the entire contents of which are each incorporated herein by reference.

VISCERAL PAIN

Visceral pain can lead to visceral reactions which can manifest themselves as, for example, contractions of the abdominal muscles. The number of contractions of the abdominal muscles occurring after a mechanical pain stimulus produced by distending the large intestine can thus be a measurement for determining visceral sensitivity to pain.

The inhibiting action of a test agent on distension-induced contractions can be tested in rats. The distension of the large intestine with an introduced balloon can be used as the stimulus; the contraction of the abdominal muscles can be measured as the response.

For example, one hour after sensitizing of the large intestine by instillation of a weak acetic acid solution, a latex balloon is introduced and inflated sequentially in a stepwise fashion to about 50-100 mbar for about 5-10 minutes. Pressure values can also be expressed as cm H₂O at 4°C (mbar X 1.01973 = cm H₂O at 4°C). During this time, the contractions of the abdominal muscles are counted. About 20 minutes after subcutaneous administration of the test agent, this measurement is repeated.

The action of the test agent is calculated as a percentage reduction in the counted contractions compared with the control (i.e., non-sensitized rats).

VISCEROMOTOR RESPONSE TO COLORECTAL DISTENSION

The ability of a treatment to reverse acetic acid-induced colonic hypersensitivity in a rodent model of irritable bowel syndrome can be assessed by the following method.

Animals

Adult male Fisher rats are housed under standard conditions. Following one week of acclimatization to the animal facility, the rats are brought to the laboratory and handled daily for another week to get used to the environment and the research associate performing the experiments.

Visceromotor Responses to Colorectal Distension (CRD)

The visceromotor behavioral response to colorectal distension was measured by counting the number of abdominal contractions recorded by a strain gauge sutured onto the abdominal musculature as described in Gunter *et al.*, *Physiol. Behav.*, 69(3): 379-82 (2000) in awake unrestrained animals. A 5 cm latex balloon catheter inserted via the anal canal into the colon is used for colorectal distensions. Constant pressure tonic distensions were performed in a graded manner (15, 30 or 60 mmHg) and are maintained for a period of 10 min and the numbers of abdominal muscle contractions were recorded to measure the level of colonic sensation. A 10 min recovery is allowed between distensions.

Acetic Acid-Induced Colonic Hypersensitivity

Acetic acid-induced colonic hypersensitivity in rats has been described by Langlois *et al.*, *Eur. J. Pharmacol.*, 318: 141-144 (1996) and Plourde *et al.*, *Am. J. Physiol.* 273: G191-G196 (1997). In the present study, a low concentration of acetic acid (1.5 ml, 0.6%) was administered intracolonicly to sensitize the colon without causing histological damage to the colonic mucosa as described in previous studies

(Gunter *et al.*, *supra*).

Testing

Drug(s) or vehicle alone are administered to the rats 30 min prior to initiation of the protocol for colorectal distension. Injection volume was 0.2mL using 100% propylene glycol as the vehicle. Three consecutive colorectal distensions at 15, 30 or 60 mmHg applied at 10-min intervals were recorded. Visceromotor responses were evaluated as the number of abdominal muscle contractions recorded during the 10-min periods of colorectal distension. Non-sensitized and sensitized uninjected control animals serve to demonstrate the lower and upper levels of response, respectively.

GASTROINTESTINAL (GI) MOTILITY MODEL

The investigation of gastrointestinal motility can be based on either the in vivo recording of mechanical or electrical events associated intestinal muscle contractions in whole animals or the activity of isolated gastrointestinal intestinal muscle preparations recorded in vitro in organ baths (see, for example, Yaun *et al.*, *Br. J. Pharmacol.*, 112(4):1095-1100 (1994), Jin *et al.*, *J. Pharm. Exp. Ther.*, 288(1): 93-97 (1999) and Venkova *et al.*, *J. Pharm. Exp. Ther.*, 300(3):1046-1052 (2002)). The in vivo recordings, especially in conscious freely moving animals, have the advantage of characterizing motility patterns and propulsive activity that are directly relevant to the motor function of the GI tract. In comparison, in vitro studies provide data about the mechanisms and site of action of agents directly affecting contractile activity and are a classic tool to distinguish between effects on the circular and /or longitudinal intestinal smooth muscle layers.

IN VIVO COLONIC CONTRACTILITY

Ambulatory telemetric motility recordings provide a suitable way to investigate intestinal motility in conscious animals during long-lasting time periods. Telemetric recording of colonic motility has been introduced to study propagating contractile activity in the unprepared colon of conscious freely moving animals. Yucatan mini-pigs, present an excellent animal model for motility investigations,

based on the anatomical and functional similarities between the gastrointestinal tract in the human and the mini-pig. To be prepared for studies of colonic motility, young mini-pigs undergo a surgical procedure to establish a permanent chronic cecal fistula.

During an experimental trial, the animals are housed in an animal facility under controlled conditions and receive a standard diet with water available ad libitum. Telemetric recording of colonic motility in a segment of proximal colon in the mini-pig is carried out for approximately one week (McRorie *et al.*, *Dig. Dis. Sci.* 43: 957-963 (1998); Kuge *et al.*, *Dig. Dis. Sci.* 47: 2651-6 (2002)). The data obtained in each recording session can be used to define the mean amplitude and the total number of propagating contractions, the number of high and low velocity propagating contractions, the number of long and short duration propagating contractions and to estimate the relative shares of each type contractions as % of total contractile activity. A summarized motility index (MI), characterizing colonic contractile activity, can be calculated using the following equation:

$$MI = \# \text{ of contractions} / 24 \text{ hr.} \times \text{area under the pressure peak } 24 \text{ hr.}$$

COLONIC MOTILITY

Female rats are administered, TNBS in ethanol or saline (control), intracolonicly. The catheter tip is positioned between 2 and 6 cm from the anal verge (n=6/group). Three days following TNBS administration, the animals are food restricted overnight and on the following morning are anesthetized with urethane and are instrumented for physiological/pharmacological experimentation.

A ventral incision is made on the ventral surface of the neck, a jugular catheter is inserted and secured with ligatures, and the skin wound is closed with suture. An intra-colonic balloon-tipped catheter fashioned from condom reservoir tip and tubing is inserted anally and positioned with the balloon at approximately 4 cm from the anal verge. Connection via 3-way stopcock to a syringe pump and pressure transducer allows for simultaneous balloon volume adjustment and pressure recording. Fine wire electrodes are inserted into the external anal sphincter (EAS) and the abdominal wall musculature to permit electromyographic (EMG) recording.

With this preparation, intra-colonic pressure, colonic motility, colonic sensory thresholds via abdominal EMG firing, and EAS firing frequency and amplitude is quantified in both control and irritated animals.

Following a control period of about 1 hour at a balloon volume of about 0.025 ml to establish baseline colonic motility and associated non-noxious viscerosomatic reflex measurements, three consecutive escalating ramps of stepwise or continuous balloon inflation are conducted. Following the completion of each volume ramp, the balloons are deflated for 30 minutes for recovery and collection of additional colonic motility measurements. EMG and colonic pressure responses to balloon inflation are measured and analyzed as sensitivities to colorectal distension (CRD). Administration of pharmacological agents is conducted in an escalating dose-response protocol and begins following the last control CRD balloon deflation.

IN VITRO

Recordings of contractile activity of isolated smooth muscle preparations can be used to study selected aspects of muscle function under conditions where the influence of "external" factors (circulating hormones etc.) is removed, while the muscle itself retains its in vivo capacity.

Studies are performed using smooth muscle strips (or whole intestinal segments) mounted vertically in organ baths with one end fixed and the other attached to isometric force transducers. The muscles are continuously bathed in modified Krebs bicarbonate buffer, maintained at 37°C and aerated with 95% O and 5% CO . The tissues are allowed to equilibrate at initial length (Li- at which tension is zero) for approximately 5 minutes, and then are gradually stretched by small force increments to optimal length (Lo - the length at which maximal active tension is generated in response to an agonist). Experiments should be performed at Lo to provide standardized spontaneous activity and pharmacological responses. The most commonly used recording procedures involve isometric transducers attached to an appropriate recording device. Mechanical responses to stimulation of enteric nerve terminals can be studied in organ baths supplied with pairs of platinum electrodes connected to a physiological electrical stimulator. Isolated smooth muscle

preparations can be used also to study length-tension relationships, which provide characteristics of the active and passive properties of the smooth muscle.

A MODEL OF INCREASED COLONIC TRANSIT

The model used in this example provides a method of determining the ability of a drug(s) to normalize accelerated colonic transit induced by water avoidance stress (WAS). This model provides a method of evaluating the effectiveness of a compound in a specific patient group of IBS sufferers where stress induced colonic motility is considered a significant contributing factor. Preliminary testing in the water avoidance stress model confirmed that there exists an association between stress and altered colonic motility. Fecal pellet output is measured by counting the total number of fecal pellets produced during 1 hour of WAS.

Animals

Adult male F-344 rats, supplied by Charles River Laboratories and weighing 270-350 g, were used to complete this study. The rats were housed 2 per cage under standard conditions. Following one to two weeks of acclimatization to the animal facility, the rats were brought to the laboratory and handled daily for another week to acclimatize them to laboratory conditions and to the research associate who performed the studies. All procedures used in this study were approved in accordance with facility standards.

Acclimatization prior to experiments

All rats underwent sham stress (1-hour in stress chamber without water) for 2-4 consecutive days before undergoing WAS (sham was performed until rats produced 0-1 pellet per hour for 2 consecutive days). At the end of the 1-hour stress period, the fecal pellets were counted and recorded.

Procedure

WAS causes an acceleration of colonic transit, which can be quantified by counting the number of fecal pellets, produced during the stress procedure. Rats were placed for 1-hour into a stress chamber onto a raised platform 7.5 cm x 7.5 cm

x 9 cm (L x W x H) in the center of a stress chamber filled with room temperature 10 water 8 cm in depth. The stress chamber was constructed from a rectangular plastic tub (40.2 x 60.2 x 31.2 cm).

SMALL INTESTINAL TRANSIT

The effect of a drug(s) on the inhibition of small intestinal transit can be evaluated using the Small Intestinal Transit Rodent Model described below.

Method

Adult male F-344 rats are used in this study. The rats are housed under standard conditions. The animals are fed a standard rodent diet and food and water are provided "ad libitum". Rats are allowed to acclimatize to the animal facility for one week prior to the transit experiments. Small intestinal transit in rats is investigated by the passage of a charcoal meal along the small intestine during a defined time period (15-min). The animals are deprived of food for 12-16 hrs prior to the experiments. Rats are given a charcoal meal (a mixture of charcoal, gum arabic, and distilled water) as a 2 mL oral gavage and are sacrificed after a 15-min test period. The distance traveled by the charcoal meal is quantified as a percent of the small intestinal length, using the following equation: $\text{Transit (\%)} = \text{cm traveled by meal} / \text{cm total small intestinal length} \times 100$.

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following embodiments are encompassed by the present invention:

1. A pharmaceutical composition comprising an $\alpha_2\delta$ subunit calcium channel modulator and an antimuscarinic in an amount sufficient to produce a synergistic effect in the treatment of functional bowel disorders.
2. The pharmaceutical composition of embodiment 1 wherein said antimuscarinic is oxybutynin in the treatment of functional bowel disorders.
3. The pharmaceutical composition of embodiment 1 wherein said $\alpha_2\delta$ subunit calcium channel modulator is a GABA analog in the treatment of functional bowel disorders.
4. The pharmaceutical composition of embodiment 3 wherein said GABA analog is gabapentin in the treatment of functional bowel disorders.
5. The pharmaceutical composition of embodiment 4 wherein said antimuscarinic is oxybutynin in the treatment of functional bowel disorders.
6. A pharmaceutical composition comprising gabapentin and oxybutynin or pharmaceutically acceptable acids, salts, esters, amides, prodrugs, or active metabolites thereof in an amount sufficient to treat functional bowel disorders.
7. The pharmaceutical composition of embodiment 6 wherein said gabapentin is present in an amount from about 600 mg to about 2400 mg, and wherein said oxybutynin is present in an amount less than about 5 mg.
8. The pharmaceutical composition of embodiment 6 wherein said gabapentin and oxybutynin are present in a ratio based on a fraction of their respective

ED₅₀ values, and wherein said ratio is from about 1:1 to about 300:1 or from about 1:1 to about 1:300, respectively.

9. A combination for the treatment of functional bowel disorders comprising amounts of gabapentin and oxybutynin or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof in a weight/weight ratio of from 1:1 to about 300:1 or from about 1:1 to about 1:300, respectively.

10. A method for treating functional bowel disorders, which comprises administering to an individual in need thereof a therapeutically effective amount of a first component that is an $\alpha_2\delta$ subunit calcium channel modulator or a pharmaceutically acceptable acid, salt, ester, amide, prodrug, or active metabolite thereof in combination with a second component that is a smooth muscle modulator or a pharmaceutically acceptable acid, salt, ester, amide, prodrug, or active metabolite thereof.

11. The method of embodiment 10, wherein the functional bowel disorder is selected from the group consisting of IBS, functional abdominal bloating, functional constipation, functional diarrhea, functional abdominal pain syndrome, and functional dyspepsia.

12. The method of embodiment 10, wherein the functional bowel disorder is IBS.

13. The method of embodiment 10, wherein the functional bowel disorder is functional abdominal bloating.

14. The method of embodiment 10, wherein the functional bowel disorder is functional constipation.

15. The method of embodiment 10, wherein the functional bowel disorder is functional diarrhea.


16. The method of embodiment 10, wherein the active agents are contained within a pharmaceutical formulation.

17. The method of embodiment 16, wherein the pharmaceutical formulation is a unit dosage form.


18. The method of embodiment 10, wherein the $\alpha_2\delta$ subunit calcium channel modulator is selected from the group consisting of:


- a. Gabapentin and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Pregabalin and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. GABA analogs as described in U.S. Pat. No. 4,024,175 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. GABA analogs as described in U.S. Pat. No. 5,563,175 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. GABA analogs as described in U.S. Patent No. 6,316,638 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. GABA analogs as described in PCT Publication No. WO 93/23383 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. GABA analogs as described in Bryans *et al.* (1998) *J. Med. Chem.* 41:1838-1845 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. GABA analogs as described in Bryans *et al.* (1999) *Med. Res. Rev.* 19:149-177 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- i. Amino acid compounds as described in U.S. Application No. 20020111338 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Cyclic amino acid compounds as disclosed in PCT Publication No. WO 99/08670 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. Cyclic amino acids (illustrated below) as disclosed in PCT Publication No. WO99/21824 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;









- l. Bicyclic amino acids (illustrated below) as disclosed in published U.S. Patent Application No. 60/160725 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and





- m. Bicyclic amino acid analogs (illustrated below) as disclosed in UK Patent Application GB 2 374 595 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.





19. The method of embodiment 10, wherein said smooth muscle modulator is selected from the group consisting of: antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors.

20. The method of embodiment 19, wherein said smooth muscle modulator is an antimuscarinic.

21. The method of embodiment 20, wherein the antimuscarinic is selected from the group consisting of:

- a. Darifenacin (Daryon[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Solifenacin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. YM-905 (solifenacin succinate) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. Solifenacin monohydrochloride or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Oxybutynin (Ditropan[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. S-Oxybutynin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. N-desethyl-oxybutynin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. Tolterodine (Detrol[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Propiverine (Detrunorm[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Propantheline bromide (Pro-Banthine[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. Hyoscyamine sulfate (Levsin[®], Cystospaz[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- l. Dicyclomine hydrochloride (Bentyl[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. Flavoxate hydrochloride (Urispas[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. d,l (racemic) 4- diethylamino-2-butynyl phenylcyclohexylglycolate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. (R)-N,N-diisopropyl-3-(2-hydroxy-5-methylphenyl)-3-phenylpropanamine L-hydrogen tartrate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. (+)-(1S,3'R)-quinuclidin-3'-yl 1-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate monosuccinate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. alpha(+)-4-(Dimethylamino)-3-methyl-1,2-diphenyl-2-butanol propionate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- r. 1-methyl-4-piperidyl diphenylpropoxyacetate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. 3"-hydroxyspiro[1"H,5"H-nortropane-8,1'-pyrrolidinium benzilate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- t. 4 amino-piperidine containing compounds as disclosed in Diouf *et al.* (2002) *Bioorg. Med. Chem. Lett.* 12: 2535-9 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- u. Pirenzepine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- v. Methocitramine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. 4-diphenylacetoxv-N-methyl piperidine methiodide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. Tropicamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- y. (2R)-N-[1-(6-aminopyridin-2-ylmethyl)piperidin-4-yl]-2-[(1R)-3,3-difluorocyclopentyl]-2-hydroxy-2-phenylacetamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- z. PNU-200577 ((R)-N, N-diisopropyl-3-(2-hydroxy-5 hydroxymethylphenyl)-3-phenylpropanamine) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- aa. KRP-197(4-(2-methylimidazolyl)-2,2-diphenylbutyramide) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- bb. Fesoterodine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- cc. SPM 7605 (the active metabolite of Fesoterodine), or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof

22. The method of embodiment 19, wherein said smooth muscle modulator is a β 3 adrenergic agonist.

23. The method of embodiment 22, wherein the β 3 adrenergic agonist is selected from the group consisting of:

- a. TT-138 and phenylethanolamine compounds or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. FR-149174 and propanolamine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. KUC-7483 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. 4'-hydroxynorephedrine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. 2-amino-1-phenylethanol compounds or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. GS 332 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. BRL-37,344 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. BRL-26830A or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. CGP 12177 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. CL 316243 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. ICI 215,001 HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. ZD 7114 HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. Pindolol or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- n. (S)-(-)-Pindolol or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. SR 59230A HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. SR 58611 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- q. YM178 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof

24. The method of embodiment 19, wherein said smooth muscle modulator is a spasmolytic.

25. The method of embodiment 24, wherein the spasmolytic is selected from the group consisting of:

- a. α - α -diphenylacetic acid-4-(N-methyl-piperidyl) esters or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Dioxazocine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. Quaternary 6,11-dihydro-dibenzo-[b,e]-thiepine-11-N-alkylnorscopine ethers or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. Quaternary salts of dibenzo[1,4]diazepinones, pyrido-[1,4]benzodiazepinones, pyrido[1,5]benzodiazepinones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Endo-8,8-dialkyl-8-azoniabicyclo (3.2.1) octane-6,7-exo-epoxy-3-alkyl-carboxylate salts or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- f. Triazinones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. Piperazino-pyrimidines or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. Aralkylamino carboxylic acids or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Aralkylamino sulfones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. Smooth muscle spasmolytic agents as disclosed in US Patent No. 6,207,852 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- j. Papaverine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

26. The method of embodiment 19, wherein said smooth muscle modulator is a neurokinin receptor antagonist.

27. The method of embodiment 26, wherein the neurokinin receptor antagonist is selected from the group consisting of:

- a. RP 67580 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. CP 96,345 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. SR 48968 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. MEN 10,627 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. L 659,877 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and

- f. TAK-637 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

28. The method of embodiment 19, wherein said smooth muscle modulator is a bradykinin receptor antagonist.

29. The method of embodiment 28, wherein the bradykinin receptor antagonist is selected from the group consisting of:

- a. des-arg¹⁰HOE 140 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. des-Arg⁹bradykinin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. NPC 349 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. NPC 567 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. HOE 140 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. MEN11270 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. Icatibant or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. FR173567 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- i. WIN 64338 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

30. The method of embodiment 19, wherein said smooth muscle modulator is a nitric oxide donor.

31. The method of embodiment 30, wherein the nitric oxide donor is selected from the group consisting of:

- a. Nitroglycerin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Sodium nitroprusside or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. FK 409 (NOR-3) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. FR 144420 (NOR-4) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. 3-morpholinocydonimine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. Linsidomine chlorohydrate ("SIN-1") or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. S-nitroso-N-acetylpenicillamine ("SNAP") or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. AZD3582 (CINOD lead compound, available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. NCX 4016 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. NCX 701 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. NCX 1022 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- l. HCT 1026 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. NCX 1015 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. NCX 950 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. NCX 1000 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. NCX 1020 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. AZD 4717 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- r. NCX 1510/NCX 1512 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. NCX 2216 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- t. NCX 4040 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- u. Nitric oxide donors as disclosed in U.S. Patent No. 5,155,137 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- v. Nitric oxide donors as disclosed in U.S. Patent No. 5,366,997 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. Nitric oxide donors as disclosed in U.S. Patent No. 5,405,919 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. Nitric oxide donors as disclosed in U.S. Patent No. 5,650,442 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- y. Nitric oxide donors as disclosed in U.S. Patent No. 5,700,830 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- z. Nitric oxide donors as disclosed in U.S. Patent No. 5,632,981 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- aa. Nitric oxide donors as disclosed in U.S. Patent No. 6,290,981 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- bb. Nitric oxide donors as disclosed in U.S. Patent No. 5,691,423 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- cc. Nitric oxide donors as disclosed in U.S. Patent No. 5,721,365 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- dd. Nitric oxide donors as disclosed in U.S. Patent No. 5,714,511 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- ee. Nitric oxide donors as disclosed in U.S. Patent No. 6,511,911 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and

- ff. Nitric oxide donors as disclosed in U.S. Patent No. 5,814,666 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.
32. The method of embodiment 10, wherein the active agents are administered on an as-needed basis.
33. The method of embodiment 16, wherein the formulation is a controlled release dosage form.
34. The method of embodiment 33, wherein the formulation is a delayed release dosage form.
35. The method of embodiment 33, wherein the formulation is a sustained release dosage form.
36. The method of embodiment 34, wherein the formulation is a sustained release dosage form.
37. The method of embodiment 35, wherein the sustained release dosage form provides drug release over a time period of from about 6 hours to about 8 hours.
38. The method of embodiment 10, wherein the active agents are administered orally.
39. The method of embodiment 15, wherein the active agents are administered orally.
40. The method of embodiment 39, wherein the pharmaceutical formulation is selected from the group consisting of tablets, capsules, caplets, solutions, suspensions, syrups, granules, beads, powders and pellets.


41. The method of embodiment 40, wherein the pharmaceutical formulation comprises a tablet.
42. The method of embodiment 40, wherein the pharmaceutical formulation comprises a capsule.
43. The method of embodiment 10, wherein the active agents are administered transmucosally.
44. The method of embodiment 43, wherein the active agents are administered sublingually.
45. The method of embodiment 43, wherein the active agents are administered buccally.
46. The method of embodiment 43, wherein the active agents are administered intranasally.
47. The method of embodiment 43, wherein the active agents are administered transurethrally.
48. The method of embodiment 43, wherein the active agents are administered rectally.
49. The method of embodiment 43, wherein the active agents are administered by inhalation.
50. The method of embodiment 10, wherein the active ingredient is administered intravesically.

51. The method of embodiment 10, wherein the active agents are administered topically.
52. The method of embodiment 10, wherein the active agents are administered transdermally.
53. The method of embodiment 10, wherein the active agents are administered parenterally.
54. The method of embodiment 10, wherein the active agents are administered intrathecally.
55. The method of embodiment 15, wherein the active agents are administered intrathecally.
56. The method of embodiment 15, wherein the active agents are administered by a route of administration selected from the group consisting of: vaginally and perivaginally.
57. The method of embodiment 56, wherein the formulation is selected from the group consisting of vaginal suppositories, creams, ointments, liquid formulations, pessaries, tampons, gels, pastes, foams and sprays.
58. A pharmaceutical formulation for treating functional bowel disorders and adapted for transmucosal drug administration, comprising a therapeutically effective amount of a first component that is an $\alpha_2\delta$ subunit calcium channel modulator or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof in combination with a second component that is a smooth muscle modulator or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, and a carrier suitable for transmucosal drug delivery buccally, sublingually, intranasally, rectally, or by inhalation.



59. The formulation of embodiment 58 wherein the functional bowel disorder is selected from the group consisting of IBS, functional abdominal bloating, functional constipation, functional diarrhea, functional abdominal pain syndrome, and functional dyspepsia.
60. The formulation of embodiment 59, wherein the functional bowel disorder is IBS.
61. The formulation of embodiment 59, wherein the functional bowel disorder is functional abdominal bloating.
62. The formulation of embodiment 59, wherein the functional bowel disorder is functional constipation.
63. The formulation of embodiment 59, wherein the functional bowel disorder is functional diarrhea.
64. The formulation of embodiment 57, comprising a solid dosage form for application to the buccal mucosa, and wherein the carrier is suitable for buccal drug delivery.
65. The formulation of embodiment 64, wherein the carrier is a hydrolyzable polymer.
66. The formulation of embodiment 64, wherein the dosage form further comprises an adhesive suitable for affixing the dosage form to the buccal mucosa.
67. The formulation of embodiment 58, comprising a dosage form for application to the sublingual mucosa, and wherein the carrier is suitable for sublingual drug delivery.

68. The formulation of embodiment 58, comprising a dosage form for application to the rectal mucosa, and wherein the carrier is suitable for rectal drug delivery.
69. The formulation of embodiment 68, comprising a rectal suppository.
70. The formulation of embodiment 58, comprising a dosage form suitable for inhalation.
71. The formulation of embodiment 70, comprising a liquid.
72. The formulation of embodiment 70, comprising a dry powder.
73. The formulation of embodiment 70, comprising an aerosol composition.
74. The formulation of embodiment 58, wherein the active agents are administered by a route of administration selected from the group consisting of: vaginally and perivaginally.
75. The formulation of embodiment 74, wherein the formulation is selected from the group consisting of vaginal suppositories, creams, ointments, liquid formulations, pessaries, tampons, gels, pastes, foams and sprays.
76. The formulation of embodiment 58, wherein the $\alpha_2\delta$ subunit calcium channel modulator is selected from the group consisting of:
- a. Gabapentin and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
 - b. Pregabalin and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;





- c. GABA analogs as described in U.S. Pat. No. 4,024,175 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. GABA analogs as described in U.S. Pat. No. 5,563,175 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. GABA analogs as described in U.S. Patent No. 6,316,638 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. GABA analogs as described in PCT Publication No. WO 93/23383 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. GABA analogs as described in Bryans *et al.* (1998) *J. Med. Chem.* 41:1838-1845 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. GABA analogs as described in Bryans *et al.* (1999) *Med. Res. Rev.* 19:149-177 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Amino acid compounds as described in U.S. Application No. 20020111338 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Cyclic amino acid compounds as disclosed in PCT Publication No. WO 99/08670 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. Cyclic amino acids (illustrated below) as disclosed in PCT Publication No. WO99/21824 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;




- l. Bicyclic amino acids (illustrated below) as disclosed in published U.S. Patent Application No. 60/160725 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and



- m. Bicyclic amino acid analogs (illustrated below) as disclosed in UK Patent Application GB 2 374 595 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.





77. The formulation of embodiment 58, wherein said smooth muscle modulator is selected from the group consisting of: antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors.

78. The formulation of embodiment 77, wherein said smooth muscle modulator is an antimuscarinic.

79. The formulation of embodiment 78, wherein the antimuscarinic is selected from the group consisting of:

- a. Darifenacin (Daryon[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Solifenacin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. YM-905 (solifenacin succinate) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- d. Solifenacin monohydrochloride or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Oxybutynin (Ditropan[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. S-Oxybutynin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. N-desethyl-oxybutynin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. tolterodine (Detrol[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Propiverine (Detrunorm[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Propantheline bromide (Pro-Banthine[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. Hyoscyamine sulfate (Levsin[®], Cystospaz[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. Dicyclomine hydrochloride (Bentyl[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. Flavoxate hydrochloride (Urispas[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. d,l (racemic) 4- diethylamino-2-butynyl phenylcyclohexylglycolate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- o. (R)-N,N-diisopropyl-3-(2-hydroxy-5-methylphenyl)-3-phenylpropanamine L-hydrogen tartrate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. (+)-(1S,3'R)-quinuclidin-3'-yl 1-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate monosuccinate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. alpha(+)-4-(Dimethylamino)-3-methyl-1,2-diphenyl-2-butanol propionate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- r. 1-methyl-4-piperidyl diphenylpropoxyacetate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. 3"-hydroxyspiro[1 "H,5"H-nortropane-8,1'-pyrrolidinium benzilate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- t. 4 amino-piperidine containing compounds as disclosed in Diouf *et al.* (2002) *Bioorg. Med. Chem. Lett.* 12: 2535-9 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- u. pirenzpine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- v. methoctramine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. 4-diphenylacetoxy-N-methyl piperidine methiodide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. tropicamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- y. (2R)-N-[1-(6-aminopyridin-2-ylmethyl)piperidin-4-yl]-2-[(1R)-3,3-difluorocyclopentyl]-2-hydroxy-2-phenylacetamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- z. PNU-200577 ((R)-N, N-diisopropyl-3-(2-hydroxy-5-hydroxymethylphenyl)-3-phenylpropanamine) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- aa. KRP-197(4-(2-methylimidazolyl)-2,2-diphenylbutyramide) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- bb. Fesoterodine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- cc. SPM 7605 (the active metabolite of Fesoterodine), or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

80. The formulation of embodiment 77, wherein said smooth muscle modulator is a $\beta 3$ adrenergic agonist.

81. The formulation of embodiment 80, wherein the $\beta 3$ adrenergic agonist is selected from the group consisting of:

- a. TT-138 and phenylethanolamine compounds or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. FR-149174 and propanolamine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. KUC-7483 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- d. 4'-hydroxynorephedrine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. 2-amino-1-phenylethanol compounds or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. GS 332 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. BRL-37,344 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. BRL-26830A or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. CGP 12177 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. CL 316243 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. ICI 215,001 HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. ZD 7114 HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. Pindolol or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. (S)-(-)-Pindolol or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. SR 59230A HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. SR 58611 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. YM178 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof

82. The formulation of embodiment 77, wherein said smooth muscle modulator is a spasmolytic.

83. The formulation of embodiment 82, wherein the spasmolytic is selected from the group consisting of:

- a. α - α -diphenylacetic acid-4-(N-methyl-piperidyl) esters or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Dioxazocine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. Quaternary 6,11-dihydro-dibenzo-[b,e]-thiepine-11-N-alkyl-norscopine ethers or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. Quaternary salts of dibenzo[1,4]diazepinones, pyrido[1,4]benzodiazepinones, pyrido[1,5]benzodiazepinones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Endo-8,8-dialkyl-8-azoniabicyclo (3.2.1) octane-6,7-exo-epoxy-3-alkyl-carboxylate salts or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. Triazinones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. Piperazino-pyrimidines or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. Aralkylamino carboxylic acids or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Aralkylamino sulfones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- j. Smooth muscle spasmolytic agents as disclosed in US Patent No. 6,207,852 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- k. Papaverine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

84. The formulation of embodiment 77, wherein said smooth muscle modulator is a neurokinin receptor antagonist.

85. The formulation of embodiment 84, wherein the neurokinin receptor antagonist is selected from the group consisting of:

- a. RP 67580 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. CP 96,345 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. SR 48968 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. MEN 10,627 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. L 659,877 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- f. TAK-637 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

86. The formulation of embodiment 77, wherein said smooth muscle modulator is a bradykinin receptor antagonist.

87. The formulation of embodiment 86, wherein the bradykinin receptor antagonist is selected from the group consisting of:

- a. des-arg¹⁰HOE 140 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- b. des-Arg⁹bradykinin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. NPC 349 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. NPC 567 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. HOE 140 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. MEN11270 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. Icatibant or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. FR173567 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- i. WIN 64338 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

88. The formulation of embodiment 77, wherein said smooth muscle modulator is a nitric oxide donor.

89. The formulation of embodiment 88, wherein the nitric oxide donor is selected from the group consisting of:

- a. Nitroglycerin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Sodium nitroprusside or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. FK 409 (NOR-3) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. FR 144420 (NOR-4) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- e. 3-morpholinosydnonimine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. Linsidomine chlorohydrate ("SIN-1") or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. S-nitroso-N-acetylpenicillamine ("SNAP") or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. AZD3582 (CINOD lead compound, available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. NCX 4016 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. NCX 701 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. NCX 1022 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. HCT 1026 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. NCX 1015 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. NCX 950 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- o. NCX 1000 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. NCX 1020 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. AZD 4717 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- r. NCX 1510/NCX 1512 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. NCX 2216 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- t. NCX 4040 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- u. Nitric oxide donors as disclosed in U.S. Patent No. 5,155,137 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- v. Nitric oxide donors as disclosed in U.S. Patent No. 5,366,997 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. Nitric oxide donors as disclosed in U.S. Patent No. 5,405,919 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. Nitric oxide donors as disclosed in U.S. Patent No. 5,650,442 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- y. Nitric oxide donors as disclosed in U.S. Patent No. 5,700,830 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- z. Nitric oxide donors as disclosed in U.S. Patent No. 5,632,981 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- aa. Nitric oxide donors as disclosed in U.S. Patent No. 6,290,981 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- bb. Nitric oxide donors as disclosed in U.S. Patent No. 5,691,423 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- cc. Nitric oxide donors as disclosed in U.S. Patent No. 5,721,365 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- dd. Nitric oxide donors as disclosed in U.S. Patent No. 5,714,511 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- ee. Nitric oxide donors as disclosed in U.S. Patent No. 6,511,911 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- ff. Nitric oxide donors as disclosed in U.S. Patent No. 5,814,666 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

90. The formulation of embodiment 58, wherein the formulation is administered to an individual suffering from a functional bowel disorder.

91. The formulation of embodiment 90 wherein the functional bowel disorder is selected from the group consisting of IBS, functional abdominal bloating, functional

constipation, functional diarrhea, functional abdominal pain syndrome, and functional dyspepsia.

92. The formulation of embodiment 91, wherein the functional bowel disorder is IBS.

93. The formulation of embodiment 91, wherein the functional bowel disorder is functional abdominal bloating.

94. The formulation of embodiment 91, wherein the functional bowel disorder is functional constipation.

95. The formulation of embodiment 91, wherein the functional bowel disorder is functional diarrhea.

96. A pharmaceutical formulation for the treatment of functional bowel disorders comprising oxybutynin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof in an amount less than about 5 mg together with a pharmaceutically acceptable adjuvant, diluent or carrier.

97. The pharmaceutical formulation of embodiment 96 wherein said oxybutynin or pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof is present in an amount of about 2.5 mg.

98. The pharmaceutical formulation of embodiment 96 wherein said oxybutynin or pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof is present in an amount of about 1.5 mg.

99. A pharmaceutical formulation for the treatment of functional bowel disorders comprising gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, oxybutynin or a pharmaceutically acceptable salt

ester, amide, prodrug, or active metabolite thereof, and at least one pharmaceutically-acceptable excipient selected from the group consisting of fillers, binders, compression aids, disintegrants, lubricants, stabilizers, and solubilizers.

100. The pharmaceutical formulation of embodiment 99 wherein said gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof is present in an amount of about 200 mg.

101. The pharmaceutical formulation of embodiment 99 wherein said oxybutynin or a pharmaceutically acceptable salt ester, amide, prodrug, or active metabolite thereof is present in an amount of about 2.5 mg.

102. The pharmaceutical formulation of embodiment 99 wherein said formulation comprises gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, oxybutynin or a pharmaceutically acceptable salt ester, amide, prodrug, or active metabolite thereof, at least one filler, at least one binder, at least one compression aid, at least one disintegrant, at least one lubricant, and at least one solubilizer.

103. The pharmaceutical formulation of embodiment 102 wherein said formulation comprises gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, oxybutynin or a pharmaceutically acceptable salt ester, amide, prodrug, or active metabolite thereof, lactose, providone, microcrystalline cellulose, croscopvidone, magnesium stearate, and water.

104. The pharmaceutical formulation of embodiment 102 wherein said formulation comprises gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, oxybutynin or a pharmaceutically acceptable salt ester, amide, prodrug, or active metabolite thereof, lactose, hydroxypropylmethylcellulose, microcrystalline cellulose, croscopvidone, magnesium stearate, and water.

128. The pharmaceutical formulation of embodiment 99 wherein said formulation is a tablet.

129. A combination product for the treatment of functional bowel disorders comprising: (A) gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof; and (B) oxybutynin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, wherein each of components (A) and (B) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

130. The combination product of embodiment 129 which comprises a pharmaceutical formulation including gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, and oxybutynin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

131. A packaged kit for a patient to use in the treatment of functional bowel disorders, comprising: a pharmaceutical formulation comprising a therapeutically effective amount of a first component that is an $\alpha_2\delta$ subunit calcium channel modulator or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof in combination with a second component that is a smooth muscle modulator or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof; a container housing the pharmaceutical formulation during storage and prior to administration; and instructions for carrying out drug administration in a manner effective to treat functional bowel disorders.

132. The packaged kit of embodiment 131, wherein the pharmaceutical formulation is an oral dosage form containing a unit dosage of a first component that is an $\alpha_2\delta$ subunit calcium channel modulator or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof in combination with a unit dosage of a

second component that is a smooth muscle modulator or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, the unit dosage being a therapeutically effective dosage for treatment of functional bowel disorders .

133. The packaged kit of embodiment 131, wherein the functional bowel disorder is selected from the group consisting of IBS, functional abdominal bloating, functional constipation, functional diarrhea, functional abdominal pain syndrome, and functional dyspepsia.

134. The packaged kit of embodiment 133, wherein the functional bowel disorder is IBS.

135. The packaged kit of embodiment 133, wherein the functional bowel disorder is functional abdominal bloating.


136. The packaged kit of embodiment 133, wherein the functional bowel disorder is functional constipation.

137. The packaged kit of embodiment 133, wherein the functional bowel disorder is functional diarrhea.


138. The packaged kit of embodiment 136, wherein the $\alpha_2\delta$ subunit calcium channel modulator is selected from the group consisting of:

- a. Gabapentin and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Pregabalin and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. GABA analogs as described in U.S. Pat. No. 4,024,175 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;


- d. GABA analogs as described in U.S. Pat. No. 5,563,175 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. GABA analogs as described in U.S. Patent No. 6,316,638 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. GABA analogs as described in PCT Publication No. WO 93/23383 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. GABA analogs as described in Bryans *et al.* (1998) *J. Med. Chem.* 41:1838-1845 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. GABA analogs as described in Bryans *et al.* (1999) *Med. Res. Rev.* 19:149-177 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Amino acid compounds as described in U.S. Application No. 20020111338 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Cyclic amino acid compounds as disclosed in PCT Publication No. WO 99/08670 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. Cyclic amino acids (illustrated below) as disclosed in PCT Publication No. WO99/21824 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;








1. Bicyclic amino acids (illustrated below) as disclosed in published U.S. Patent Application No. 60/160725 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and



- m. Bicyclic amino acid analogs (illustrated below) as disclosed in UK Patent Application GB 2 374 595 and acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.





139. The packaged kit of embodiment 136, wherein said smooth muscle modulator is selected from the group consisting of: antimuscarinics, β 3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors.

140. The packaged kit of embodiment 139, wherein said smooth muscle modulator is an antimuscarinic.

141. The packaged kit of embodiment 140, wherein the antimuscarinic is selected from the group consisting of:

- a. Darifenacin (Daryon[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Solifenacin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. YM-905 (solifenacin succinate) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. Solifenacin monohydrochloride or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Oxybutynin (Ditropan[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. S-Oxybutynin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. N-desethyl-oxybutynin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. tolterodine (Detrol[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- i. Propiverine (Detrunorm[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Propantheline bromide (Pro-Banthine[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. Hyoscyamine sulfate (Levsin[®], Cystospaz[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. Dicyclomine hydrochloride (Bentyl[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. Flavoxate hydrochloride (Urispas[®]) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. d,l (racemic) 4- diethylamino-2-butynyl phenylcyclohexylglycolate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. (R)-N,N-diisopropyl-3-(2-hydroxy-5-methylphenyl)-3-phenylpropanamine L-hydrogen tartrate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. (+)-(1S,3'R)-quinuclidin-3'-yl 1-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate monosuccinate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. alpha(+)-4-(Dimethylamino)-3-methyl-1,2-diphenyl-2-butanol propionate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- r. 1-methyl-4-piperidyl diphenylpropoxyacetate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. 3"-hydroxyspiro[1"H,5"H-nortropane-8,1'-pyrrolidinium benzilate or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- t. 4 amino-piperidine containing compounds as disclosed in Diouf *et al.* (2002) *Bioorg. Med. Chem. Lett.* 12: 2535-9 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- u. pirenzipine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- v. methoctramine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. 4-diphenylacetoxy-N-methyl piperidine methiodide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. tropicamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- y. (2R)-N-[1-(6-aminopyridin-2-ylmethyl)piperidin-4-yl]-2-[(1R)-3,3-difluorocyclopentyl]-2-hydroxy-2-phenylacetamide or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- z. PNU-200577 ((R)-N, N-diisopropyl-3-(2-hydroxy-5-hydroxymethylphenyl)-3-phenylpropanamine) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- aa. KRP-197(4-(2-methylimidazolyl)-2,2-diphenylbutyramide) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- bb. Fesoterodine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- cc. SPM 7605 (the active metabolite of Fesoterodine), or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

142. The packaged kit of embodiment 139, wherein said smooth muscle modulator is a $\beta 3$ adrenergic agonist.

143. The packaged kit of embodiment 142, wherein the $\beta 3$ adrenergic agonist is selected from the group consisting of:

- a. TT-138 and phenylethanolamine compounds or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. FR-149174 and propanolamine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. KUC-7483 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. 4'-hydroxynorephedrine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. 2-amino-1-phenylethanol compounds or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. GS 332 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. BRL-37,344 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. BRL-26830A or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- i. CGP 12177 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. CL 316243 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. ICI 215,001 HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. ZD 7114 HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. Pindolol or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. (S)-(-)-Pindolol or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. SR 59230A HCl or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. SR 58611 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. YM178 or acids, salts, esters, amides, prodrugs, active metabolites, and other derivatives thereof

144. The packaged kit of embodiment 139, wherein said smooth muscle modulator is a spasmolytic.

145. The packaged kit of embodiment 144, wherein the spasmolytic is selected from the group consisting of:

- a. α - α -diphenylacetic acid-4-(N-methyl-piperidyl) esters or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Dioxazocine derivatives or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- c. Quaternary 6,11-dihydro-dibenzo-[b,e]-thiepine-11-N-alkyl-norscopine ethers or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. Quaternary salts of dibenzo[1,4]diazepinones, pyrido-[1,4]benzodiazepinones, pyrido[1,5]benzodiazepinones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. Endo-8,8-dialkyl-8-azoniabicyclo (3.2.1) octane-6,7-exo-epoxy-3-alkyl-carboxylate salts or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. Triazinones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. Piperazino-pyrimidines or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. Aralkylamino carboxylic acids or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. Aralkylamino sulfones or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. Smooth muscle spasmolytic agents as disclosed in US Patent No. 6,207,852 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- k. Papaverine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

146. The packaged kit of embodiment 139, wherein said smooth muscle modulator is a neurokinin receptor antagonist.

147. The packaged kit of embodiment 146, wherein the neurokinin receptor antagonist is selected from the group consisting of:

- a. RP 67580 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. CP 96,345 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. SR 48968 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. MEN 10,627 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. L 659,877 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- f. TAK-637 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

148. The packaged kit of embodiment 139, wherein said smooth muscle modulator is a bradykinin receptor antagonist.

149. The packaged kit of embodiment 148, wherein the bradykinin receptor antagonist is selected from the group consisting of:

- a. des-arg¹⁰HOE 140 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. des-Arg⁹bradykinin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. NPC 349 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. NPC 567 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. HOE 140 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. MEN11270 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- g. Icatibant or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- h. FR173567 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- i. WIN 64338 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

150. The packaged kit of embodiment 139, wherein said smooth muscle modulator is a nitric oxide donor.

151. The packaged kit of embodiment 150, wherein the nitric oxide donor is selected from the group consisting of:

- a. Nitroglycerin or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- b. Sodium nitroprusside or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- c. FK 409 (NOR-3) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- d. FR 144420 (NOR-4) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- e. 3-morpholinisydnnonimine or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- f. Linsidomine chlorohydrate ("SIN-1") or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- g. S-nitroso-N-acetylpenicillamine ("SNAP") or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- h. AZD3582 (CINOD lead compound, available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- i. NCX 4016 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- j. NCX 701 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- k. NCX 1022 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- l. HCT 1026 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- m. NCX 1015 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- n. NCX 950 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- o. NCX 1000 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- p. NCX 1020 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- q. AZD 4717 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- r. NCX 1510/NCX 1512 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- s. NCX 2216 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- t. NCX 4040 (available from NicOx S.A.) or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- u. Nitric oxide donors as disclosed in U.S. Patent No. 5,155,137 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- v. Nitric oxide donors as disclosed in U.S. Patent No. 5,366,997 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- w. Nitric oxide donors as disclosed in U.S. Patent No. 5,405,919 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- x. Nitric oxide donors as disclosed in U.S. Patent No. 5,650,442 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- y. Nitric oxide donors as disclosed in U.S. Patent No. 5,700,830 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- z. Nitric oxide donors as disclosed in U.S. Patent No. 5,632,981 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- aa. Nitric oxide donors as disclosed in U.S. Patent No. 6,290,981 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;

- bb. Nitric oxide donors as disclosed in U.S. Patent No. 5,691,423 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- cc. Nitric oxide donors as disclosed in U.S. Patent No. 5,721,365 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- dd. Nitric oxide donors as disclosed in U.S. Patent No. 5,714,511 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof;
- ee. Nitric oxide donors as disclosed in U.S. Patent No. 6,511,911 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof; and
- ff. Nitric oxide donors as disclosed in U.S. Patent No. 5,814,666 or acids, salts, enantiomers, analogs, esters, amides, prodrugs, active metabolites, and derivatives thereof.

152. A packaged kit for a patient to use in the treatment of functional bowel disorders, comprising: (a) a pharmaceutical formulation including gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; and (b) a pharmaceutical formulation including oxybutynin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier, which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

153. The packaged kit of embodiment 152, wherein components (a) and (b) are suitable for sequential, separate and/or simultaneous use in the treatment of functional bowel disorders.

154. A packaged kit for a patient to use in the treatment of functional bowel disorders, comprising: (a) a pharmaceutical formulation including gabapentin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; and (b) a pharmaceutical formulation including oxybutynin or a pharmaceutically acceptable salt, ester, amide, prodrug, or active metabolite thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier, which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

155. The packaged kit of embodiment 154, wherein components (a) and (b) are suitable for sequential, separate and/or simultaneous use in the treatment of functional bowel disorders.

156. A method for processing an insurance claim under an insurance policy for treatment of functional bowel disorders using an $\alpha_2\delta$ subunit calcium channel modulator and an antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof, wherein said $\alpha_2\delta$ subunit calcium channel modulator and antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof are administered sequentially or concurrently in different compositions, comprising:

- a. Receiving notification that treatment using said $\alpha_2\delta$ subunit calcium channel modulator and said antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs or active metabolites thereof will be performed or notification of a prescription;
- b. Determining whether said treatment using said $\alpha_2\delta$ subunit calcium channel modulator and said antimuscarinic or pharmaceutically acceptable

salts, esters, amides, prodrugs or active metabolites is covered under said insurance policy; and

- c. Processing said claim for treatment of said functional bowel disorders using said $\alpha_2\delta$ subunit calcium channel modulator and said antimuscarinic or pharmaceutically acceptable salts, esters, amides, prodrugs, or active metabolites thereof, including payment, reimbursement, or application against a deductible.

157. The method of embodiment 156, wherein said $\alpha_2\delta$ subunit calcium channel modulator is gabapentin.

158. The method of embodiment 156, wherein said antimuscarinic is oxybutynin.

CLAIMS

What is claimed is:

1. A method for treating functional bowel disorders which comprises administering to an individual in need thereof a therapeutically effective amount of a first component that is an $\alpha_2\delta$ subunit calcium channel modulator in combination with a second component that is a smooth muscle modulator, wherein:
 - a. said functional bowel disorder is selected from the group consisting of: irritable bowel syndrome, functional bloating, functional constipation, functional diarrhea, functional abdominal pain syndrome, and functional dyspepsia;
 - b. said $\alpha_2\delta$ subunit calcium channel modulator is selected from the group consisting of: Gabapentin; pregabalin; GABA analogs as described in U.S. Patent Nos. 4,024,175; 5,563,175; 6,316,638; GABA analogs as described in PCT Publication No. WO 93/23383; GABA analogs as described in Bryans *et al.* (1998) *J. Med. Chem.* 41:1838-1845 and Bryans *et al.* (1999) *Med. Res. Rev.* 19:149-177; amino acid compounds as described in U.S. Application No. 20020111338; cyclic amino acid compounds as disclosed in PCT Publication Nos. WO 99/08670 and WO 99/21824; bicyclic amino acids as disclosed in published U.S. Patent Application No. 60/160725; bicyclic amino acid analogs as disclosed in UK Patent Application GB 2 374 595; and derivatives and analogs thereof; and
 - c. said smooth muscle modulator is selected from the group consisting of: Darifenacin (Daryon[®]); solifenacin; YM-905 (solifenacin succinate); solifenacin monohydrochloride; Oxybutynin (Ditropan[®]); S-Oxybutynin; N-desethyl-oxybutynin; tolterodine (Detrol[®]); Propiverine (Detrunorm[®]); Propantheline bromide (Pro-Banthine[®]); Hyoscyamine sulfate (Levsin[®], Cystospaz[®]); Dicyclomine hydrochloride (Bentyl[®]); Flavoxate hydrochloride (Urispas[®]); d,l (racemic) 4- diethylamino-2-butynyl phenylcyclohexylglycolate; (R)-N,N-diisopropyl-3-(2-hydroxy-

5-methylphenyl)-3-phenylpropanamine L-hydrogen tartrate; (+)-(1S,3'R)-quinuclidin-3'-yl 1-phenyl-1,2,3,4-tetrahydroisoquinoline-2-carboxylate monosuccinate; alpha(+)-4-(Dimethylamino)-3-methyl-1,2-diphenyl-2-butanol proprionate; 1-methyl-4-piperidyl diphenylpropoxyacetate; 3"-hydroxyspiro[1"H,5"H-nortropane-8,1'-pyrrolidinium benzilate; 4 amino-piperidine containing compounds as disclosed in Diouf *et al.* (2002) *Bioorg. Med. Chem. Lett.* 12: 2535-9; pirenzipine; methocramine; 4-diphenylacetoxy-N-methyl piperidine methiodide; tropicamide; (2R)-N-[1-(6-aminopyridin-2-ylmethyl)piperidin-4-yl]-2-[(1R)-3,3-difluorocyclopentyl]-2-hydroxy-2-phenylacetamide; PNU-200577 ((R)-N, N-diisopropyl-3-(2-hydroxy-5-hydroxymethylphenyl)-3-phenylpropanamine); KRP-197(4-(2-methylimidazolyl)-2,2-diphenylbutyramide); Fesoterodine; SPM 7605 (the active metabolite of Fesoterodine); TT-138 and phenylethanolamine compounds; FR-149174 and propanolamine derivatives; KUC-7483; 4'-hydroxynorephedrine derivatives; 2-amino-1-phenylethanol compounds; GS 332; BRL-37,344; BRL-26830A; CGP 12177; CL 316243; ICI 215,001 HCl; ZD 7114 HCl; Pindolol; (S)-(-)-Pindolol; SR 59230A HCl; SR 58611; YM178; α - α -diphenylacetic acid-4-(N-methyl-piperidyl) esters; Dioxazocine derivatives; Quaternary 6,11-dihydro-dibenzo-[b,e]-thiepine-11-N-alkyl norscopine ethers; Quaternary salts of dibenzo[1,4]diazepinones, pyrido-[1,4]benzodiazepinones, pyrido[1,5]benzodiazepinones; Endo-8,8-dialkyl-8-azoniabicyclo (3.2.1) octane-6,7-exo-epoxy-3-alkyl-carboxylate salts; Triazinones; Piperazino-pyrimidines; Aralkylamino carboxylic acids; Aralkylamino sulfones; RP 67580; CP 96,345; SR 48968; MEN 10,627; L 659,877; L 659,877; TAK-637; des-arg¹⁰HOE 140; des-Arg⁹bradykinin; NPC 349; NPC 567; HOE 140; MEN11270; Icatibant; FR173567; WIN 64338; Nitroglycerin; Sodium nitroprusside; FK 409 (NOR-3); FR 144420 (NOR-4); 3-morpholinosydnonimine; Linsidomine chlorohydrate ("SIN-1"); S-nitroso-N-acetylpenicillamine ("SNAP"); AZD3582; NCX

4016; NCX 701; NCX 1022; HCT 1026; NCX 1015; NCX 950; NCX 1000; NCX 1020; AZD 4717; NCX 1510/NCX 1512; NCX 2216; and NCX 4040.

2. A pharmaceutical composition for treating functional bowel disorders comprising gabapentin and oxybutynin wherein said gabapentin is present in an amount from about 600 mg to about 2400 mg, and wherein said oxybutynin is present in an amount less than about 5 mg.

3. A method for treating irritable bowel syndrome which comprises administering to an individual in need thereof a therapeutically effective amount of composition comprising gabapentin and oxybutynin.

METHODS FOR TREATING FUNCTIONAL BOWEL DISORDERS USING $\alpha_2\delta$
SUBUNIT CALCIUM CHANNEL MODULATORS WITH SMOOTH MUSCLE
MODULATORS

ABSTRACT OF THE DISCLOSURE

A method is provided for using $\alpha_2\delta$ subunit calcium channel modulators or other compounds that interact with the $\alpha_2\delta$ calcium channel subunit in combination with one or more compounds with smooth muscle modulatory effects to treat functional bowel disorders in patients in need of treatment. According to the present invention, $\alpha_2\delta$ subunit calcium channel modulators include GABA analogs including gabapentin and pregabalin, fused bicyclic or tricyclic amino acid analogs of gabapentin, and amino acid compounds. Compounds with smooth muscle modulatory effects include antimuscarinics, β_3 adrenergic agonists, spasmolytics, neurokinin receptor antagonists, bradykinin receptor antagonists, and nitric oxide donors.